

Primarily About Primaries

Allan Borodin, Omer Lev, Nisarg Shah and Tyrone Strangway

Abstract

Much of the social choice literature examines *direct* voting systems, in which voters submit their ranked preferences over candidates and a voting rule picks a winner. Real-world elections and decision-making processes are often more complex and involve multiple stages. For instance, one popular voting system filters candidates through *primaries*: first, voters affiliated with each political party vote over candidates of their own party and the voting rule picks a candidate from each party, which then compete in a general election.

We present a model to analyze such multi-stage elections, and conduct the first quantitative comparison (to the best of our knowledge) of the direct and primary voting systems in terms of the quality of the elected candidate. Our main theoretical result is that every voting rule is guaranteed to perform almost as well (i.e., within a constant factor) under the primary system as under the direct system. Surprisingly, the converse does not hold: we show settings in which there exist voting rules that perform significantly better under the primary system than under the direct system. Using simulations, we see that plurality benefits significantly from using a primary system over a direct one, while Condorcet-consistent rules do not.

1 Introduction

Thomas Jefferson, like many of the US constitution’s authors, believed that political parties and factions are a bad thing (see also Hamilton et al. [18]). This view stemmed from a long history of British and English political history, in which prison sentences and executions were possible outcomes in the battle between factions for supremacy at the Royal court [32]. However, both in Britain and in the United States, once their respective legislative assemblies gained political force, parties turned out to be quite unavoidable. Even Jefferson had to start his own party, which ended up quite successful, and was able to vanquish the opposing party from political existence [35].

Fast forwarding to today, political parties have become the bedrock of parliamentary politics throughout the world. In particular, one of political parties’ main roles – if not the most important (especially in presidential systems) – is to select the candidates which are voted for by the general public. The mechanisms by which parties make this selection are varied, and they have evolved significantly throughout the past 150 years. But in the past few decades there has been a marked shift by parties throughout the world towards increasing the ability of individual party (or unaffiliated) members to influence the outcome, and in some cases, to be the only element to determine party candidates [12]. In particular, US parties have changed their election methods since the 1970s to focus the selection of presidential, congressional and state-wide candidates on popular support by party members via primaries [9].

Despite this long and established role of parties in whittling down the candidate field in elections, the treatment of a parties’ role in elections within the multiagent systems community has been quite limited. While various candidate manipulation attacks have been investigated (e.g., Sybil attacks [11]), and there is recent research into parties as a collection of similar minded candidates (e.g., in gerrymandering, across different districts), the role of parties in removing candidates has not been explored.

The focus of this paper is the *primary* voting system, in which each party’s electorate selects a winner from among the party’s candidates, and among these primary winners,

an ultimate election winner is selected by the general public. Our goal is to compare this system to the *direct* voting system, in which all voters directly vote over all candidates.

1.1 Our Results

Our contribution is twofold. First, we formulate a model which allows a quantitative comparison of the two voting systems. Our model is a spatial model of voting in which voters and candidates lie in an underlying multi-dimensional space, and voter preferences are single-peaked. This allows us to compare each candidate’s social utility in terms of its total distance to the voters. We make the evaluation metric formal using the notion of *distortion* advocated by a recent line of research [28, 6]. Our results focus on 2 parties, each selecting a single candidate, with both candidates presented to the general voting public.

Second, we use this model to present a comparison of the direct and primary voting systems. In particular, we show that no voting rule performs much worse (by more than a constant factor) under the primary system than under the direct system. While the converse holds in some cases, we show settings in which it does not, and there exist voting rules that perform much better under the primary system than under the direct system.

It is important to note that while we write of parties, voters and elections, this multi-step model applies to a variety of decision-making processes by agents. An organization selecting an “employee of the month” may ask each unit to nominate a single candidate, and then choose from amongst them. A city may ask its regional subdivisions to assess which roads require urgent fixing, and then the city council decides from these options where to invest its efforts. Fundamentally, in many cases where the potential number of options is huge, it is common to use subdivisions to cull the options and present only a few of them for discussion and vote. In such cases, our multi-stage model is apt.

2 Related Work

The analysis of regular, direct elections is long and varied, both in the social sciences and in AI [7]. In our particular setting, the voters are located in a metric space, with their preferences related to their distance from candidates. Such settings have been widely investigated in the social science literature since the work of Downs [14], recently summarized by Schofield [30]. In particular, we focus on the idea of distortion in such elections, a topic broached by Procaccia and Rosenschein [28] for voters with a utility function, but investigated in the context of voters in metric spaces in a series of papers [2, 1, 33] for most common voting rules. Feldman et al. [17] explored such a setting for strategyproof mechanisms. Micha and Shah [24] consider the problem of predicting the election outcome under a given voting rule using sampled votes. This can be alternatively viewed as the problem of choosing a desirable candidate for the entire electorate when only a subset of them vote. In our metric setting, Lemma 4 shows that if only a constant fraction of the electorate votes, the distortion of the chosen candidate with respect to the entire electorate increases by at most a constant factor. In contrast, in their non-metric setting, they derive a stark negative result even when the votes of all but one voter are available. Finally, Bachrach et al. [3] examine a related concept, comparing how district based elections differ from proportional ones. They measure how much a district election outcome can differ from the proportional outcome, a term they refer to as the *Misrepresentation Ratio*.

Discussing changes to the set of candidates has been mainly focused on two paths of research. Strategic candidates, investigation of which began with the work of Dutta et al. [15] – followed by Dutta et al. [16] discussing strategic candidacy in tournaments, and recently further explored by Brill and Conitzer [8], Polukarov et al. [27] and others – deals

mainly with finding equilibria. The other is the addition and removal of candidates, as a form of control manipulation, which was studied by Bartholdi III et al. [4]; see the summaries by Brandt et al. [7] and Rothe [29].

Investigating parties' selection methods and their effect on the election has mostly been done in the social sciences. Kenig [23] details the range of selection methods parties use, and there has been significant focus on more democratic methods for leader selection [13], which seems to be a general trend in many Western countries [12]. There is also significant literature on particular party elections in various countries, such as Britain [21], Belgium [34], Israel [20], and many others. Naturally, the most widely examined country is the US, in which political parties have been a fixture of political life since its early days [35]. The most recent extensive summary of research on it is due to Cohen et al. [9], who try to explain how party power-brokers influence the party membership vote. Norpoth [26] uses primary data to predict election results, and notably Sides et al. [31] show that primary voters are very similar to "regular" voters. In computational fields, recent interest in proxy voting [10, 22], in which voters give other agents the ability to vote for them, may be related to how modern parties are viewed and analyzed. Finally, Harrenstein et al. [19] looks at the complexity of finding Nash equilibrium for the strategic selection of party nominees.

3 Model

For $k \in \mathbb{N}$, define $[k] = \{1, \dots, k\}$. Let $V = [n]$ denote a set of n voters, and A denote a set of m candidates. We assume that voters and candidates lie in an underlying metric space $M = (S, d)$, where S is a set of points and d is a distance function satisfying the triangle inequality and symmetry. More precisely, there exists an *embedding* $\rho : V \cup A \rightarrow S$ mapping each voter and candidate to a point in S . For a set $X \subseteq V \cup A$, we slightly abuse the notation and let $\rho(X) = \{\rho(x) : x \in X\}$. Also, for $x, x' \in V \cup A$, we often use $d(x, x')$ instead of $d(\rho(x), \rho(x'))$ for notational convenience.

In this work, we assume that voters and candidates additionally have an affiliation with a political party. Specifically, we study a setting with two parties¹, denoted -1 and 1 . The *party affiliation* function $\pi : V \cup A \rightarrow \{-1, 1\}$ maps each voter and candidate to the party they are affiliated with. For $p \in \{-1, 1\}$, let $V_p = \pi^{-1}(p) \cap V$, $A_p = \pi^{-1}(p) \cap A$, $n_p = |V_p|$, and $m_p = |A_p|$. We require $n_p \geq 1$ for each $p \in \{-1, 1\}$. Our main result (Theorem 2) holds even if there are independent voters not affiliated with either party.

Collectively, an *instance* is the tuple $I = (V, A, M, \rho, \pi)$. Given I , our goal is to find a candidate $a \in A$ as the winner. The *social cost* of a is its total distance to the voters, denoted $C^I(a) = \sum_{i \in V} d(i, a)$. For party $p \in \{-1, 1\}$, let $C_p^I(a) = \sum_{i \in V_p} d(i, a)$. Hence $C^I(a) = C_{-1}^I(a) + C_1^I(a)$. We also use for $X \subseteq V$, $C_X^I(a) = \sum_{i \in X} d(i, a)$. Given an instance I , we would like to choose a candidate $a_{\text{OPT}} \in \arg \min_{a \in A} C^I(a)$ that minimizes the social cost. We shall drop the instance from superscripts if it is clear from the context.

However, we do not observe the full instance. Specifically, we do not know the underlying metric M or the embedding function ρ . Instead, each voter $i \in N$ submits a *vote*, which is a ranking (strict total order) \succ_i over the candidates in A by their distance to the voter. Specifically, for all $i \in N$ and $a, b \in A$, $a \succ_i b \Rightarrow d(i, a) \leq d(i, b)$. The voter is allowed to break ties between equidistant candidates arbitrarily. The *vote profile* $\vec{\succ}^I = (\succ_1, \dots, \succ_n)$ is the collection of votes. Given an instance I , its corresponding *election* $E^I = (V, A, \vec{\succ}^I, \pi)$ contains all observable information.

In the families of instances that we consider, we fix the number of candidates m and let the number of voters n to be arbitrarily large. This choice is justified because in many

¹The model can be extended to multiple parties in a reasonably straightforward manner, but for simplicity's sake, we shall focus on only 2 parties.

typical elections (e.g., political ones), voters significantly outnumber candidates. Let $\mathcal{I}_{m,\mathcal{M}}^\alpha$ be the family of instances satisfying the following conditions.

- Each party has at least an α fraction of the voters affiliated with it, i.e., $n_p \geq \alpha \cdot n$ for each $p \in \{-1, 1\}$. Note that $\alpha \in [0, 0.5]$: $\alpha = 0.5$ is the strictest (exactly half of the voters are affiliated with each party), while $\alpha = 0$ imposes no conditions; in the latter case, we omit the superscript α .
- The number of candidates is at most m .

In particular, we shall focus on a few cases of \mathcal{M} :

$\mathcal{M} = \star$ This allows M to be an arbitrary metric space.

$\mathcal{M} = \mathbb{R}^k$ The metric space should be $M = (\mathbb{R}^k, d)$, where d is the standard Euclidean distance.

$\mathcal{M} = \mathbf{sep}\text{-}\mathbb{R}^k$ This means the embedding ρ must be such that $\rho(V_{-1} \cup A_{-1})$ and $\rho(V_1 \cup A_1)$ are linearly separable.² In this case we shall take the metric to be $M = (\mathbb{R}^k, d)$ with d as the standard Euclidean distance. In plain words, the voters and candidates affiliated with each party reside in a certain part of the metric space, separate from those affiliated with the other party. In a single dimension, this means there exists a threshold on the line such that voters and candidates affiliated with one party lie to the left of it, while those affiliated with the other party lie to the right. Note that this is the only choice of \mathcal{M} that restricts the embedding ρ based on party affiliation π .

These families of instances are related by the following relation. For all k ,

$$\mathcal{I}_{m,\mathbf{sep}\text{-}\mathbb{R}^k}^\alpha \subset \begin{matrix} \mathcal{I}_{m,\mathbb{R}^k}^\alpha \\ \mathcal{I}_{m,\mathbf{sep}\text{-}\mathbb{R}^{k+1}}^\alpha \end{matrix} \subset \mathcal{I}_{m,\mathbb{R}^{k+1}}^\alpha \subset \mathcal{I}_{m,\star}^\alpha$$

3.1 Voting Rules and Distortion

A voting rule f takes an election as input, and returns a winning candidate from A . We say that the cost-approximation of f on instance I is

$$\phi(f, I) = \frac{C^I(f(E^I))}{\min_{a \in A} C^I(a)},$$

and given a family of instances \mathcal{I} , the distortion of f with respect to \mathcal{I} is

$$\phi_{\mathcal{I}}(f) = \sup_{I \in \mathcal{I}} \phi(f, I).$$

Since distortion is a worst-case notion, we have that when $\mathcal{I} \subseteq \mathcal{I}'$, $\phi_{\mathcal{I}}(f) \leq \phi_{\mathcal{I}'}(f)$ for every voting rule f .

Standard voting rules choose the winning candidate independently of party affiliations. These include rules such as plurality, Borda count, k -approval, veto, and STV. We refer readers to the book by Brandt et al. [7] for their definitions. We call a voting rule *affiliation-independent* if $f(E) = f(E')$ when elections E and E' differ only in their party affiliation functions. Since an affiliation-independent voting rule f ignores party affiliations, we have $\phi_{\mathcal{I}_{m,\mathbf{sep}\text{-}\mathbb{R}^k}^\alpha}^\alpha(f) = \phi_{\mathcal{I}_{m,\mathbb{R}^k}^\alpha}^\alpha(f)$. All of the above-mentioned rules, in addition to being affiliation-independent, share the property of being *unanimous*, i.e., they return candidate a when a is the top choice of all voters.

²Two sets of points are linearly separable if the interiors of their convex hulls are disjoint.

3.2 Stages and Primaries

Given an affiliation-independent voting rule f , voting systems with primaries employ a specific process to choose the winner, essentially resulting in a different voting rule \hat{f} that operates on a given election $E = (V, A, \vec{\succ}, \pi)$ as follows:

1. First, it creates two *primary elections*: for $p \in \{-1, 1\}$, define $E_p = (V_p, A_p, \vec{\succ}_p, \pi_p)$, where $\vec{\succ}_p$ denotes the preferences of voters in V_p over candidates in A_p , and $\pi_p : V_p \rightarrow \{p\}$ is a constant function.
2. Next, it computes the winning candidate in each primary election (primary winner) using rule f : for $p \in \{-1, 1\}$, let $a_p^* = f(E_p)$.
3. Finally, let $E_g = (V, \{a_{-1}^*, a_1^*\}, \vec{\succ}_g, \pi)$ be the *general election*, where $\vec{\succ}_g$ denotes the preferences of all voters over the two primary winners. The winning candidate is $\hat{f}(E) = \text{maj}(E_g)$, which is what most voting rules become when dealing with only 2 candidates.³ Here, maj is the majority rule, which, given two candidates, picks the one that a majority of voters prefer; our results are independent of its tie-breaking.

This resembles systems employed by the main US, Canadian and other countries' parties, in which a party's members vote on their party's candidates to select a winner of their primary. In other systems, such selection could be a multi-stage process.

Given an affiliation-independent voting rule f , the goal of this paper is to compare its performance under the *direct system*, in which f is applied on the given election directly, to its performance under the *primary system*, in which \hat{f} is applied on the given election instead. Formally, given a family of instances \mathcal{I} and an affiliation-independent voting rule f , we wish to compare $\phi_{\mathcal{I}}(f)$ and $\phi_{\mathcal{I}}(\hat{f})$ (henceforth, the distortion of f with respect to \mathcal{I} under the direct and the primary systems, respectively).

4 Small Primaries Are Terrible

Recall that in a family of instances $\mathcal{I}_{m, \mathcal{M}}^\alpha$, we require that at least α fraction of voters be affiliated with each party, i.e., $n_p \geq \alpha n$ for each $p \in \{-1, 1\}$. In other words, each primary election must have at least αn voters.

We first show that when a primary election can have very few voters ($\alpha = 0$), every reasonable voting rule has an unbounded distortion in the primary system, even with respect to our most stringent family of instances $\mathcal{I}_{m, \text{sep-}\mathbb{R}}$.

Theorem 1. *For $m \geq 3$, $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}}(\hat{f}) = \infty$ for every affiliation-independent unanimous voting rule f .*

Proof. Consider an instance $I \in \mathcal{I}_{m, \text{sep-}\mathbb{R}}$ in which voter 1 is located at 0 and affiliated with party -1 , while the remaining $n - 1$ voters are located at 1 and affiliated with party 1. All m candidates are affiliated with party -1 : one is at 0, and the rest are at 1.

The candidate a^* at 0 becomes the primary winner of party -1 , and trivially becomes the overall winner. Its social cost is $C(a^*) = n - 1$. In contrast, an optimal candidate a_{OPT} at 1 has social cost $C(a_{OPT}) = 1$. Hence, $\frac{C(a^*)}{C(a_{OPT})} = n - 1$. Since the number of voters n is unbounded, $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}}(\hat{f}) = \infty$. \square

³If one of the parties has no affiliated candidates, then the primary winner of the other party becomes the overall winner. In a setting with more than 2 parties, or where each party nominates several candidates, the general election can use f to determine the outcome (or use some other voting process).

Theorem 1 continues to hold even if we require that at least a constant fraction of candidates be affiliated with each party: we could simply move a constant fraction of the candidates from 1 to 3, and the proof would still go through.

On the other hand, if we require that at least a *constant fraction of voters* be affiliated with each party, the result changes dramatically.

5 Large Primaries Are Never Much Worse

For every affiliation-independent voting rule f , we bound the distortion of \widehat{f} in terms of the distortion of f for every instance. Note that this is stronger than comparing the worst-case distortions of f and \widehat{f} over a family of instances.

Given an instance $I = (V, A, M, \rho, \pi)$ and party $p \in \{-1, 1\}$, we say that $I_p = (V_p, A_p, M, \rho_p, \pi_p)$ is the *primary instance* of party p , where ρ_p and π_p are restrictions of ρ and π to $V_p \cup A_p$. The primary election E_p of party p is precisely the election corresponding to instance I_p .

Theorem 2. *Let $I = (V, A, M, \rho, \pi)$ be an instance. For $p \in \{-1, 1\}$, let I_p be the primary instance of party p , and $n_p = |V_p| \geq \alpha n$. Then,*

$$\phi(\widehat{f}, I) \leq 3 \cdot \frac{1 - \alpha + \max(\phi(f, I_{-1}), \phi(f, I_1))}{\alpha}.$$

Further, for a socially optimal candidate $a_{OPT} \in \arg \min_{a \in A} C^I(a)$, we have a bound depending only on the distortion of the primary election of its party:

$$\phi(\widehat{f}, I) \leq 3 \cdot \frac{1 - \alpha + \phi(f, I_{\pi(a_{OPT})})}{\alpha}.$$

For each family of instances \mathcal{I} that we study, it holds that for every instance $I \in \mathcal{I}$, both its primary instances, if seen as direct elections, are also in \mathcal{I} (since the party division has no effect on the direct election distortion). Hence, we can convert the instance-wise comparison to a worst-case comparison.

Corollary 3. *For every $\alpha > 0$, $k \in \mathbb{N}$, family of instances $\mathcal{I} \in \{\mathcal{I}_{m, \star}, \mathcal{I}_{m, \mathbb{R}^k}, \mathcal{I}_{m, sep-\mathbb{R}^k}\}$, and affiliation-independent voting rule f ,*

$$\phi_{\mathcal{I}}(\widehat{f}) \leq 3 \cdot \frac{1 - \alpha + \phi_{\mathcal{I}}(f)}{\alpha}.$$

Note that $\phi_{\mathcal{I}}(f) \geq 1$ by definition. Hence, we can write $\phi_{\mathcal{I}}(\widehat{f}) \leq \frac{6}{\alpha} \cdot \phi_{\mathcal{I}}(f)$. In other words, for every affiliation-independent voting rule f , its distortion under the primary system is at most a constant times bigger than its distortion under the direct system, with respect to every family of instances that we consider.

To prove Theorem 2, we need two lemmas. The first lemma shows that if the distortion of a rule f for a primary instance I_p is low, then the corresponding primary winner a_p^* is nearly as good as any candidate in A_p for the overall election as well. The intuition is that when voters in $V \setminus V_p$ drive up the social cost of a_p^* (i.e., when they are far from a^*), they must do so for every candidate in A_p .

Lemma 4. *Let a_p^* denote the primary winner of party p . Then*

$$C(a_p^*) \leq \frac{1 - \alpha + \phi(f, I_p)}{\alpha} \cdot \min_{a \in A_p} C(a).$$

Proof. Let $\theta = \phi(f, I_p)$ (hence $\theta \geq 1$). Fix an arbitrary $a \in A_p$. Then, $C_{V_p}(a_p^*) \leq \theta \cdot C_{V_p}(a)$. Now,

$$\begin{aligned} C(a_p^*) &= C_{V_p}(a_p^*) + C_{V \setminus V_p}(a_p^*) \\ &\leq \theta \cdot C_{V_p}(a) + C_{V \setminus V_p}(a) + (n - n_p) \cdot d(a, a_p^*) \\ &\leq \theta \cdot C(a) + (n - n_p) \cdot d(a_p^*, a), \end{aligned} \quad (1)$$

where the second transition follows due to the triangle inequality. We also have $d(a_p^*, a) \leq d(a_p^*, i) + d(i, a)$ for any $i \in V_p$. Thus,

$$\begin{aligned} d(a_p^*, a) &\leq \frac{C_{V_p}(a_p^*) + C_{V_p}(a)}{n_p} \\ &\leq \frac{1 + \theta}{n_p} \cdot C_{V_p}(a) \leq \frac{1 + \theta}{n_p} \cdot C(a). \end{aligned} \quad (2)$$

Substituting Equation (2) into Equation (1), and using the fact that $\frac{n - n_p}{n_p} \leq \frac{1 - \alpha}{\alpha}$, we get the desired result. \square

Our next lemma shows that the primary winner that wins the general election is not much worse than the primary winner that loses the general election.

Lemma 5. *Let a_{-1}^* and a_1^* be the two primary winners, $a^* \in \{a_{-1}^*, a_1^*\}$ be the winner of the general election, and $\hat{a} \in \{a_{-1}^*, a_1^*\} \setminus \{a^*\}$. Then,*

$$C(a^*) \leq 3 \cdot C(\hat{a}).$$

Proof. Because a^* wins the general election by a majority vote, there must exist $X \subseteq V$, $|X| \geq \frac{n}{2}$ such that $d(i, a^*) \leq d(i, \hat{a})$ for every $i \in X$. Combining with the triangle inequality $d(a^*, \hat{a}) \leq d(a^*, i) + d(i, \hat{a})$, we get $d(i, \hat{a}) \geq \frac{d(a^*, \hat{a})}{2}$ for every $i \in X$. Hence,

$$C(\hat{a}) \geq \frac{n}{2} \cdot \frac{d(a^*, \hat{a})}{2} \Rightarrow d(a^*, \hat{a}) \leq \frac{4}{n} \cdot C(\hat{a}). \quad (3)$$

Now, we have

$$\begin{aligned} C(a^*) &= \sum_{i \in X} d(i, a^*) + \sum_{i \in V \setminus X} d(i, a^*) \\ &\leq \sum_{i \in X} d(i, \hat{a}) + \sum_{i \in V \setminus X} (d(i, \hat{a}) + d(a^*, \hat{a})) \\ &\leq C(\hat{a}) + \frac{n}{2} \cdot d(a^*, \hat{a}), \end{aligned} \quad (4)$$

where the second transition follows because $d(i, a^*) \leq d(i, \hat{a})$ for $i \in X$, and $d(i, a^*) \leq d(i, \hat{a}) + d(a^*, \hat{a})$ due to the triangle inequality. Substituting Equation (3) into Equation (4), we get the desired result. \square

We are now ready to prove our main result.

Proof of Theorem 2. Recall that a_{-1}^* and a_1^* are the primary winners. Let $p \in \{-1, 1\}$ be such that a_p^* is the winner of the general election. Let $a_{OPT} \in \arg \min_{a \in A} C(a)$ be a socially optimal candidate. We consider three cases.

- *Case 1:* $a_{OPT} \in A_p$. That is, the optimal candidate and the winner are affiliated with the same party. In this case, Lemma 4 yields

$$\phi(\hat{f}, I) \leq \frac{1 - \alpha + \phi(f, I_p)}{\alpha} = \frac{1 - \alpha + \phi(f, I_{\pi(a_{OPT})})}{\alpha}.$$

- *Case 2:* $a_{OPT} \in A_{-p}$, $a_{OPT} = a_{-p}^*$. That is, the optimal candidate and the winner are affiliated with different parties, and the optimal candidate is a primary winner. In this case, Lemma 5 yields

$$\phi(\widehat{f}, I) \leq 3.$$

- *Case 3:* $a_{OPT} \in A_{-p}$, $a_{OPT} \neq a_{-p}^*$. That is, the optimal candidate is affiliated with a party different than that of the winner, and is not a primary winner. In this case, we use both Lemmas 4 and 5 to derive

$$\begin{aligned} \phi(\widehat{f}, I) &= \frac{C(a_p^*)}{C(a_{OPT})} = \frac{C(a_p^*)}{C(a_{-p}^*)} \cdot \frac{C(a_{-p}^*)}{C(a_{OPT})} \\ &\leq 3 \cdot \frac{1 - \alpha + \phi(f, I_{-p})}{\alpha} \\ &= 3 \cdot \frac{1 - \alpha + \phi(f, I_{\pi(a_{OPT})})}{\alpha}. \end{aligned}$$

Thus, in each case, we have the desired approximation. \square

Note that our proof of Theorem 2 does not preclude the existence of *independent voters*. Specifically, we can allow the party affiliation function π to map a subset of voters V_0 to a neutral choice (say 0), have these voters only vote in the general election and not in either primary election under the primary system, and the proof of Theorem 2 would still go through. Additionally, we can also relax the restriction that all voters vote in the general election. That is, we can assume that a subset of voters $V_g \subseteq V$ vote in the general election under the primary system, assume $|V_g| \geq \gamma n$, and a version of Theorem 2 in which the constant 3 in the bound is replaced by $\frac{4-\gamma}{\gamma}$ would hold. This requires a generalization of Lemma 5, omitted due to space constraints. Finally, note we do not use the assumption that both parties use the same voting rule f in their primaries. The theorem extends easily to allow the use of different voting rules, with the distortion under the primary system still being bounded in terms of the maximum of the distortions in the two primary elections.

6 Large Primaries Are Not Better Without Party Separability

While we showed in the previous section that a voting rule does not perform much worse under the primary system than under the direct system, we now show that it does not perform any better either, at least in the worst case over all instances with at most m candidates. The result continues to hold even if we require each party to have at least a constant fraction of the voters.

Note, this result is weaker than Theorem 2 because it is a worst-case comparison instead of an instance-wise comparison. However, it still applies to all voting rules f . It applies to any metric that does not require separability of parties, in particular to $\mathcal{I}_{m,\star}$ and $\mathcal{I}_{m,\mathbb{R}^k}$.

Theorem 6. *For $\alpha \in [0, 0.5]$, $k \in \mathbb{N}$, $\mathcal{M} \in \{\star, \mathbb{R}^k\}$ (i.e. when the metric space does not require party separability), and affiliation-independent voting rule f , we have $\phi_{\mathcal{I}_{m,\mathcal{M}}^\alpha}(\widehat{f}) \geq \phi_{\mathcal{I}_{m,\mathcal{M}}^\alpha}(f)$.*

Proof in appendix.

Our proof actually establishes a slightly stronger result. Instead of showing $\phi_{\mathcal{I}_{m,\mathcal{M}}^\alpha}(\widehat{f}) \geq \phi_{\mathcal{I}_{m,\mathcal{M}}^\alpha}(f)$, we actually show $\phi_{\mathcal{I}_{m,\mathcal{M}}^{0.5}}(\widehat{f}) \geq \phi_{\mathcal{I}_{m,\mathcal{M}}^\alpha}(f)$.

7 Separability and Its Advantages

The analysis for $\mathcal{I}_{m,\text{sep-}\mathbb{R}^k}$ is not so straightforward. In the proof of Theorem 6, we co-located the new voters affiliated with party 1 and a_{OPT} affiliated with party -1 . This was allowed because non-separable metrics like \star and \mathbb{R}^k place no constraints on the embedding.

With $\mathcal{I}_{m,\text{sep-}\mathbb{R}^k}^\alpha$, we need the voters and candidates affiliated with one party to be separated from those affiliated with the other. Hence, this operation of putting all of one party's voters at the location of a_{OPT} belonging to another party would be allowed only if, in the original instance I , a_{OPT} is on the boundary of the convex hull of $\rho(V \cup A)$. While this is not the case for all instances, we only need this in at least one *worst-case instance* for f , i.e., for at least one $I \in \mathcal{I}_{m,\text{sep-}\mathbb{R}^k}^\alpha$ with $\phi(f, I) = \phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}^k}^\alpha}^\alpha(f)$. Equation (5) would then yield the desired result. More generally, it is sufficient if, given any $\epsilon > 0$, we can find an instance I such that $\phi(f, I) \geq \phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}^k}^\alpha}^\alpha(f) - \epsilon$ and a_{OPT} is at distance at most ϵ from the boundary of the convex hull of $\rho(V \cup A)$.

Interestingly, [2] show that this is indeed the case for plurality and Borda count (see the proof of their Theorem 4). Thus, we have the following.

Proposition 7. *Let f be plurality or Borda count. Then, $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^{0.5}}(\hat{f}) \geq \phi_{\mathcal{I}_{m,\star}}(f)$.*

However, known worst cases for the Copeland rule [2] and STV [33] do not satisfy this requirement. It is unknown if these rules admit a different worst case that satisfies it.

This raises an important question. *Does Proposition 7 hold for all affiliation-independent voting rules?* We shall shortly answer this *negatively*.

More precisely, we construct an affiliation-independent voting rule f such that $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha}^\alpha(\hat{f}) \ll \phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha}^\alpha(f)$ for every $\alpha > 0$. That is, with large primaries, f performs much better under the primary system than under the direct system, when voters and candidates are embedded on a line and the separability condition is imposed.

Note that instances in $\mathcal{I}_{m,\text{sep-}\mathbb{R}}$ are highly structured. For instance, it is known that when voters and candidates are embedded on a line, there always exists a weak Condorcet winner [5], and selecting such a candidate results in a distortion of at most 3 [2]. Hence, we have $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}}(f) = 3$ for every Condorcet-consistent, affiliation-independent voting rule f .⁴

Our aim in this section is to construct an affiliation-independent voting rule f_{fail} that with respect to $\mathcal{I}_{m,\text{sep-}\mathbb{R}}$ has an unbounded distortion in the direct system, but at most a constant distortion in the primary system.

Definition 1. Let f_{fail} be an affiliation-independent voting rule that operates on election $E = (V, A, \succ)$ as follows. Let $A = \{a_1, \dots, a_m\}$, and $t = (m + 1)/2$.

- *Special Case:* If $m \geq 9$, m is odd, $n \geq m^2$, and \succ has the following structure, then return a_1 .

1. For voter 1, $a_1 \succ_1 \dots \succ_1 a_m$.
2. For voter 2, $a_m \succ_2 \dots \succ_2 a_1$.
3. For voter 3, a_{t-1} is the most preferred, and $a_{m-2} \succ_3 a_1 \succ_3 a_{m-1} \succ_3 a_m$.
4. For voter 4, a_{t+1} is the most preferred, and $a_3 \succ_4 a_m \succ_4 a_2 \succ_4 a_1$.
5. For $j \in [m - 2]$, for voter $i = 4 + (2j - 1)$, $a_{j+1} \succ_i a_{j+2} \succ_i a_j$, and for voter $i' = 4 + 2j$, $a_{j+1} \succ_{i'} a_j \succ_{i'} a_{j+2}$.

⁴[2] also proved that no affiliation-independent (deterministic) voting rule can have distortion better than 3, even with respect to $\mathcal{I}_{m,\text{sep-}\mathbb{R}}$.

6. For every other voter v , a_t is the most preferred.

- If E does not fall under the special case, then apply any Condorcet consistent voting rule (e.g., Copeland’s rule).

Theorem 8. For $m \geq 9$ and constant $\alpha \in (0, 0.5]$, $\phi_{\mathcal{I}_{m, \text{sep-R}}^\alpha}(\widehat{f}_{\text{fail}})$ is upper bounded by a constant, whereas $\phi_{\mathcal{I}_{m, \text{sep-R}}^\alpha}(f_{\text{fail}})$ is unbounded.

Proof is in the appendix.

8 Using Simulations to Go Beyond Worst Case

So far we compared the distortion of a voting rule under the direct and primary systems, taken in the *worst case* over a family of instances. In practice, such worst-case instances may not arise naturally. In this section, we investigate the distortion of a voting rule under the direct and primary systems, in the *average case* over simulated instances. Our simulations focus on the two-stage, two-party primary process which has been main setting in this paper. We generate the simulated instances by varying a number of parameters; to keep the number of simulations reasonable, when varying one parameter, we use default values of the others⁵:

Total voters The number of voters n : default = 500, range = 100 to 2100 in increments of 200.

Total candidates The number of candidates m : default = 50, range = 10 to 210 in increments of 20.

Independent voters The percentage of voters who are independent (i.e., do not vote in any party primary): default = 0%, range = 0% to 90% in increments of 10%.

Independent candidates The percentage of candidates who are independent (i.e., are not a candidate in any party primary): default = 0%, range = 0% to 90% in increments of 10%.

Party voter balance The percentage of voters who are affiliated with party -1 : default = 50%,⁶ range = 10% to 90% in increments of 10%.

Party candidate balance The percentage of candidates who are affiliated with party -1 : default = 50%,⁷ range = 10% to 90% in increments of 10%.

Metric space dimension The dimension k of the Euclidean metric space $[0, 1]^k$: default = 4, range = $\{1, 4, 7, 10\}$.

For a given combination of the parameter values, we generate random instances as follows. First, we place a set V of n voters at uniformly random locations in $[0, 1]^k$. Next, if the ratio of the number of voters in the two parties is supposed to be $x : (1 - x)$, we find a hyperplane dividing voters into this $x : (1 - x)$ ratio. We do not set voter affiliations yet. Next, if the ratio of the number of candidates in the two parties is supposed to be $x : (1 - x)$,

⁵As we observe later, the dimension of the metric space had perhaps the most significant effect on the distortion. To verify that this was not an artifact of the default values of the other parameters, we varied the dimension along with every other parameter. Our figures, however, are generated with the default value for the dimension.

⁶When there are no independent voters in these simulations, by default we split voters equally between the two parties.

⁷When there are no independent candidates in these simulations, by default we split the candidates equally between the two parties.

then we place candidates uniformly at the appropriate side of the hyperplane. Finally, if $x\%$ of the voters (resp. candidates) are supposed to be independent, then we choose $x\%$ of the voters (resp. candidates) — rounded down — remove them from the respective sets, and distribute them at uniformly random locations in $[0, 1]^k$.

Once the locations of the voters and candidates are fixed, we create two instances. In one instance (called “split”), we assign $V_{-1} \cup A_{-1}$ to party -1 , and assign $V_1 \cup A_1$ to party 1 . In this instance, we have party separability. In the other instance (called “random”), we assign $|V_{-1}|$ voter and $|A_{-1}|$ candidates chosen uniformly at random to party -1 , and among the rest, assign $|V_1|$ voter and $|A_1|$ candidates chosen uniformly at random to party 1 . In this instance, we do not have party separability. This allows us to compare the effect of party separability on the distortion. We run five voting rules — plurality, Borda, STV, maximin, and Copeland — on both instances under the direct and primary systems, and measure the distortion (for primary systems with independent candidates, we use the specified voting rule in both rounds). Note that the distortion of the direct system would be identical for party separable and random instances because the two instances only differ in party affiliations. Thus, for each rule, we obtain three numbers: Direct, Primary-split, and Primary-random. For each combination of parameter values, we repeat this 1000 times and take the average distortion.

	Primary is better	Direct is better	No significant difference
plurality split	183	7	18
plurality random	203	0	5
STV split	8	178	22
STV random	152	13	48
Borda split	2	202	4
Borda random	8	179	21
maximin split	0	207	1
maximin random	0	208	0
Copeland split	0	207	1
Copeland random	0	208	0

Table 1: The table shows the number of settings (out of 208) in which each of primary and direct systems leads to a lower average distortion than the other. Statistical significance is measured using a paired t-test with $p = 0.05$.

We now present our results comparing the primary and direct systems. Our experiments result in 208 settings (combination of parameter values). For each setting, we compare the average primary distortion and the average direct distortion of each voting rule under each party affiliation model (split or random), and evaluate which system results in a better average distortion. For statistical significance, we use the paired t-test with $p = 0.05$. The results are presented in Table 1.

Without party separability (i.e. in the random case), our theoretical results indicate that primary is no better than direct in the worst case (Theorem 6). While this is also true in our experimentally generated average cases for Borda, Copeland, and maximin, we see that for plurality and STV, primary almost always outperforms direct in the average case.

With party separability, direct outperforms primary for all voting rules except plurality. While our theoretical result shows that for plurality direct also outperforms primary in the worst case (Proposition 7), we see that this is not true in the average case. Interestingly, for STV, party separability significantly affects which system works better.

We also wanted to know how large where the differences between the two systems, so we looked at the difference between the average distortion under the primary system and the




Figure 1: These histograms show the difference between the average distortion under the primary system and that under the direct system.

average distortion under the direct system. The results for plurality, STV, and maximin are given in Figure 1. The figures for Borda and Copeland are omitted because they were similar to the figures for maximin.

The results are quite varied. In some cases where primary mostly outperforms direct (e.g. in plurality-split or plurality-random), primary sometimes outperforms direct by a large margin whereas direct only outperforms primary by a small margin. But we see that the opposite is also true (e.g. in maximin-split). Hence, neither system seems to have a significant advantage over the other in terms of the difference in average distortions.

9 Discussion

Our paper initiates the novel quantitative study of multi-stage elections (and their comparison to single-stage elections), but leaves plenty to explore. Some directions are fairly straightforward extensions of our results. The most straightforward question is to tighten our bounds. There is also the question of explaining the trends we observe in the average case, which sometimes differ from our worst-case results. A next step would be to study realistic distributional models of voter preferences and candidate positions in the political spectrum, and analyze their effect on distortion.

Other extensions are seemingly more involved. Extending our framework to more than two parties requires the use of a ranked voting rule in the general election, which may significantly affect the analysis. Interestingly, such an extension would also incorporate independent candidates because one can imagine an independent candidate to be a party of their own. Examining the use of multiple and different voting rules as Narodytska and Walsh [25] do for two-step voting (though without candidate elimination between stages) is an enticing direction. For example, in a multi-party direct system, we may use plurality, whereas in the primary system, the parties may use STV. It is also reasonable to consider that each party has its own voting rule. It would be interesting as well to examine party manipulation techniques in primary systems. Similarly, it is reasonable to believe that candidates may also strategically shift, to some extent, their location following the primaries, to make themselves more appealing to the general electorate.

References

- [1] Elliot Anshelevich and John Postl. Randomized social choice functions under metric preferences. *Journal of Artificial Intelligence Research*, 58(1):797–827, January 2017.
- [2] Elliot Anshelevich, Onkar Bhardwaj, and John Postl. Approximating optimal social choice under metric preferences. In *Proceedings of the 29th AAAI Conference on Artificial Intelligence (AAAI)*, pages 777–783, Austin, Texas, January 2015.
- [3] Yoram Bachrach, Omer Lev, Yoad Lewenberg, and Yair Zick. Misrepresentation in district voting. In *Proceedings of the 25th International Joint Conference on Artificial Intelligence (IJCAI)*, pages 81–87, New York City, New York, July 2016.
- [4] John J. Bartholdi III, Craig A. Tovey, and Michael A. Trick. How hard is it to control an election? *Mathematical and Computer Modelling*, 16(8-9):27–40, August-September 1992.
- [5] Duncan Black. On the rationale of group decision-making. *Journal of Political Economy*, 56:23–34, 1948.
- [6] C. Boutilier, I. Caragiannis, S. Haber, T. Lu, A. D. Procaccia, and O. Sheffet. Optimal social choice functions: A utilitarian view. *Artificial Intelligence*, 227:190–213, 2015.
- [7] Felix Brandt, Vincent Conitzer, Ulle Endriss, Jérôme Lang, and Ariel D. Procaccia, editors. *Handbook of Computational Social Choice*. Cambridge University Press, March 2016.
- [8] Markus Brill and Vincent Conitzer. Strategic voting and strategic candidacy. In *Proceedings of the 29th AAAI Conference on Artificial Intelligence (AAAI)*, pages 819–826, Austin, Texas, January 2015.
- [9] Marty Cohen, David Karol, Hans Noel, and John Zaller. *The Party Decides: Presidential Nominations Before and After Reform*. University of Chicago Press, 2008.
- [10] Gal Cohensius, Shie Mannor, Reshef Meir, Eli Meïrom, and Ariel Orda. Proxy voting for better outcomes. In *Proceedings of the 16th International Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, pages 858–866, São-Paulo, Brazil, May 2017.
- [11] Vincent Conitzer and Makoto Yokoo. Using mechanism design to prevent false-name manipulations. *AI Magazine*, 31(4):65–77, 2010.
- [12] William P. Cross and André Blais. Who selects the party leader? *Party Politics*, 18(2):127–150, January 2011.
- [13] William P. Cross and Richard S. Katz, editors. *The Challenges of Intra-Party Democracy*. Oxford University Press, 2013.
- [14] Anthony Downs. *An Economic Theory of Democracy*. Harper and Row, 1957.
- [15] Bhaskar Dutta, Matthew O. Jackson, and Michel Le Breton. Strategic candidacy and voting procedures. *Econometrica*, 69(4):1013–1037, July 2001.
- [16] Bhaskar Dutta, Matthew O. Jackson, and Michel Le Breton. Voting by successive elimination and strategic candidacy. *Journal of Economic Theory*, 103(1):190–218, March 2002.

- [17] Michal Feldman, Amos Fiat, and Iddan Golomb. On voting and facility location. In *Proceedings of the 17th ACM Conference on Economics and Computation (EC)*, pages 269–286, Maastricht, The Netherlands, July 2016.
- [18] Alexander Hamilton, James Madison, and John Jay. *The Federalist Papers*. The Independent Journal, 1787.
- [19] Paul Harrenstein, Grzegorz Lisowski, Ramanujan Sridharan, and Paolo Turrini. A hotelling-downs framework for party nominees. In *Proceedings of the 20th International Joint Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, pages 593–601, London, United Kingdom, May 2021.
- [20] Reuven Y. Hazan. The 1996 intra-party elections in Israel: Adopting party primaries. *Electoral Studies*, 16(1):95–103, March 1997.
- [21] Richard Jobson and Mark Wickham-Jones. Grippled by the past: Nostalgia and the 2010 labour party leadership contest. *British Politics*, 5(4):525–548, December 2010.
- [22] Anson Kahng, Simon Mackenzie, and Ariel D. Procaccia. Liquid democracy: An algorithmic perspective. In *Proceedings of the 32nd Conference on Artificial Intelligence (AAAI)*, pages 1095–1102, New Orleans, Louisiana, February 2018.
- [23] Ofer Kenig. Classifying party leaders’ selection methods in parliamentary democracies. *Journal of Elections, Public Opinion and Parties*, 19(4):433–447, October 2009.
- [24] Micha and Nisarg Shah. Can we predict the election outcome from sampled votes? In *Proc. of 34th AAAI Conference on Artificial Intelligence*, pages 2176–2183, New York City, United States of America, February 2020.
- [25] Nina Narodytska and Toby Walsh. Manipulating two stage voting rules. In *Proceedings of the 12th International Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, pages 423–430, St. Paul, Minnesota, May 2013.
- [26] Helmut Norpoth. From primary to general election: A forecast of the presidential vote. *PS: Political Science & Politics*, 37(4):737–740, October 2004.
- [27] Maria Polukarov, Svetlana Obraztsova, Zinovi Rabinovich, Alexander Kruglyi, and Nicholas R. Jennings. Convergence to equilibria in strategic candidacy. In *Proceedings of the 24th International Conference on Artificial Intelligence (IJCAI)*, pages 624–630, Buenos Aires, Argentina, July 2015.
- [28] Ariel D. Procaccia and Jeffrey S. Rosenschein. The distortion of cardinal preferences in voting. In *Proceedings of the 10th International Workshop on Cooperative Information Agents (CIA)*, pages 317–331, Edinburgh, Great Britain, September 2006.
- [29] Jörg Rothe, editor. *Economics and Computation*. Springer, 2015.
- [30] Norman Schofield. *The Spatial Model of Politics*. Number 95 in Routledge Frontiers of Political Economy. Routledge, 2008.
- [31] John Sides, Chris Tausanovitch, Lynn Vavreck, and Christopher Warshaw. On the representativeness of primary electorates. *British Journal of Political Science*, pages 1–9, March 2018.
- [32] Brendan Simms. *Three Victories and a Defeat: The Rise and Fall of the First British Empire, 1714-1783*. Allen Lane, 2007.

- [33] Piotr Skowron and Edith Elkind. Social choice under metric preferences: Scoring rules and STV. In *Proceedings of the 31st AAAI Conference on Artificial Intelligence (AAAI)*, pages 706–712, San Francisco, California, February 2017.
- [34] Bram Wauters. Explaining participation in intra-party elections: Evidence from belgian political parties. *Party Politics*, 16(2):237–259, September 2010.
- [35] Sean Wilentz. *The Rise of American Democracy*. Norton, 2005.

Allan Borodin
 University of Toronto
 Toronto, Ontario, Canada
 Email: bor@cs.toronto.edu

Omer Lev
 Ben-Gurion University
 Beersheba, Israel
 Email: omerlev@bgu.ac.il

Nisarg Shah
 University of Toronto
 Toronto, Ontario, Canada
 Email: nisarg@cs.toronto.edu

Tyrone Strangway
 Ben-Gurion University
 Beersheba, Israel
 Email: strangwa@bgu.ac.il

A Proof of Theorem 6

Recall Theorem 6 states:

For $\alpha \in [0, 0.5]$, $k \in \mathbb{N}$, $\mathcal{M} \in \{\star, \mathbb{R}^k\}$ (i.e. when the metric space does not require party separability), and affiliation-independent voting rule f , we have $\phi_{\mathcal{I}_{m, \mathcal{M}}^\alpha}(\hat{f}) \geq \phi_{\mathcal{I}_{m, \mathcal{M}}^\alpha}(f)$.

Proof. We shall denote $\mathcal{I}_{m, \mathcal{M}}^\alpha$ as \mathcal{I} . We want to show that for every instance $I \in \mathcal{I}$, there exists an instance $I' \in \mathcal{I}$ such that $\phi(\hat{f}, I') \geq \phi(f, I)$. This would imply the desired result.

Fix an instance $I = (V, A, M, \rho, \pi) \in \mathcal{I}$. Let $a_{OPT} \in A$ denote an optimal candidate in I , and $a^* = f(E^I)$. Note that $\phi(f, I) = \frac{C^I(a^*)}{C^I(a_{OPT})}$. Construct instance $I' = (V', A, M, \rho', \pi')$ as follows.

- Let $V' = V \cup \tilde{V}$, where \tilde{V} is a new set of voters and $|\tilde{V}| = |V|$.
- Let $\rho'(x) = \rho(x)$ for all $x \in V \cup A$, and $\rho(x) = \rho(a_{OPT})$ for all $x \in \tilde{V}$. That is, ρ' matches ρ for existing voters and candidates, and the new voters are co-located with a_{OPT} .
- Let $\pi'(x) = -1$ for all $x \in V \cup A$, and $\pi'(x) = 1$ for all $x \in \tilde{V}$. That is, all existing voters and candidates are affiliated with party -1 , while all new voters are affiliated with party 1 .

First, let us check that $I' \in \mathcal{I}$. Since I has m candidates, so does I' . Further, in I' , we have $|V'_{-1}| = |V'_1| = |V'|/2$, which satisfies the constraint corresponding to every $\alpha \in [0, 0.5]$. Hence, we have $I' \in \mathcal{I}$.

Let us apply \widehat{f} on I' . One of its primary instances, I'_{-1} , is precisely I . Hence, the primary winner of party -1 is $f(I'_{-1}) = f(I) = a^*$. Because there are no candidates affiliated with party 1 , a^* becomes the overall winner.⁸

Next, $C^{I'}(a^*) \geq C^I(a^*)$ because $V \subset V'$. Also, $C^{I'}(a_{OPT}) = C^I(a_{OPT})$ because a_{OPT} has zero distance to all voters in $V' \setminus V$. Together, they yield

$$\phi(\widehat{f}, I') = \frac{C^{I'}(a^*)}{C^{I'}(a_{OPT})} \geq \frac{C^I(a^*)}{C^I(a_{OPT})} = \phi(f, I), \quad (5)$$

as desired. \square

B Proof of Theorem 8

Recall, Theorem 8 states:

For $m \geq 9$ and constant $\alpha \in (0, 0.5]$, $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}^\alpha}(\widehat{f}_{\text{fail}})$ is upper bounded by a constant, whereas $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}^\alpha}(f_{\text{fail}})$ is unbounded.

And that f_{fail} in an affiliation-independent voting rule that operates on election $E = (V, A, \succ)$ as follows. Let $A = \{a_1, \dots, a_m\}$, and $t = (m + 1)/2$.

- *Special Case:* If $m \geq 9$, m is odd, $n \geq m^2$, and \succ has the following structure, then return a_1 .
 1. For voter 1, $a_1 \succ_1 \dots \succ_1 a_m$.
 2. For voter 2, $a_m \succ_2 \dots \succ_2 a_1$.
 3. For voter 3, a_{t-1} is the most preferred, and $a_{m-2} \succ_3 a_1 \succ_3 a_{m-1} \succ_3 a_m$.
 4. For voter 4, a_{t+1} is the most preferred, and $a_3 \succ_4 a_m \succ_4 a_2 \succ_4 a_1$.
 5. For $j \in [m - 2]$, for voter $i = 4 + (2j - 1)$, $a_{j+1} \succ_i a_{j+2} \succ_i a_j$, and for voter $i' = 4 + 2j$, $a_{j+1} \succ_{i'} a_j \succ_{i'} a_{j+2}$.
 6. For every other voter v , a_t is the most preferred.
- If E does not fall under the special case, then apply any Condorcet consistent voting rule (e.g., Copeland's rule).

We will now prove the theorem. Note that m being odd ensures that t is an integer, and $m \geq 9$ ensures that $a_1, a_3, a_{t-1}, a_t, a_{t+1}, a_{m-2}$, and a_m are all distinct candidates. The significance of $n \geq m^2$ will be clear later.

We will now establish that a worst-case instance of f_{fail} falls under the special case; for this instance, we need to show that a_t is socially optimal; that f_{fail} returns a_1 on this instance; and most importantly, that the structure of \succ ensures that the optimal candidate a_t is sufficiently far from both the leftmost and the rightmost candidates.

We prove this last fact in the following lemma.

⁸Even if we require each party to have at least one affiliated candidate, the proof essentially continues to hold. In this case, we can add one candidate affiliated with party 1 that is located sufficiently far from all the voters, ensuring that a^* still becomes the overall winner. This would show $\phi_{\mathcal{I}_{m+1, \mathcal{M}}^\alpha}(\widehat{f}) \geq \phi_{\mathcal{I}_{m, \mathcal{M}}^\alpha}(f)$ because instance I' may now have $m + 1$ candidates.

Lemma 9. Let $I = (V, A, M, \rho, \pi) \in \mathcal{I}_{m, \text{sep-}\mathbb{R}}$ be an instance for which the corresponding election E^I falls under the special case of f_{fail} . Then the following holds.

1. Either $\rho(a_1) \leq \dots \leq \rho(a_m)$, or $\rho(a_1) \geq \dots \geq \rho(a_m)$, or $|\rho(A)| = 2$.
2. If $|\rho(A)| \neq 2$, $\min\{d(a_t, a_1), d(a_t, a_m)\} \geq \frac{d(a_1, a_m)}{4}$.

Proof. Since voter 1 ranks a_m last and preferences are single peaked on the line, a_m is at one edge of the candidate ordering. Similarly, since voter 2 ranks a_1 last, candidate a_1 is also at the edge of the candidate ordering (i.e., $\rho(a_1) = \max_{a \in A} \rho(a)$ or $\rho(a_1) = \min_{a \in A} \rho(a)$) and $\rho(a_m) = \max_{a \in A} \rho(a)$ or $\rho(a_m) = \min_{a \in A} \rho(a)$. If $\rho(a_1) = \rho(a_m)$, this means voters 1 and 2 are located in an equal distance from all candidates (which means all candidates are located in the same location, or some are at some distance from voters 1 and 2, and the rest are at the same distance in the other direction from these voters).

Assume $|\rho(A)| > 2$ (this also means $\rho(a_1) \neq \rho(a_m)$ and $\rho(v_1) \neq \rho(v_2)$), we wish to show the order of candidates is as voter 1 ordered them, i.e., $\rho(a_1) \leq \dots \leq \rho(a_m)$ or $\rho(a_1) \geq \dots \geq \rho(a_m)$. If voter 1 is further away from all candidates (i.e. if $\rho(a_1) = \max_{a \in A} \rho(a)$, then $\rho(v_1) > \rho(a_1)$; if $\rho(a_1) = \min_{a \in A} \rho(a)$, then $\rho(v_1) < \rho(a_1)$), the ordering of the candidates is as voter 1 orders them. Otherwise, let ℓ be the smallest index such that $\rho(a_1) \neq \rho(a_\ell)$, then $\rho(v_1)$ may be between $\rho(a_1)$ and $\rho(a_\ell)$. If $d(v_1, a_1) < d(v_1, a_\ell)$, once again, the ordering of candidates is as voter 1 ordered them. If $d(v_1, a_1) = d(v_1, a_\ell)$, for any $\ell' > \ell$, $\rho(a_{\ell'}) \neq \rho(a_1)$, as that contradicts voter 2's vote ($a_{\ell'} \succ_2 a_\ell \succ_2 a_1$). Therefore, $\rho(\ell')$ is either at $\rho(\ell)$, or further away from $\rho(a_1)$, meaning that candidates locations are ordered in the order voter 1's ordered them.

For the second condition, we will show that $d(a_t, a_1) \geq \frac{d(a_1, a_m)}{4}$. By symmetry, we also obtain $d(a_t, a_m) \geq \frac{d(a_1, a_m)}{4}$. Assume $|\rho(A)| \neq 2$, and without loss of generality, let $\rho(a_1) \leq \dots \leq \rho(a_m)$ from the first condition. We show that either $\rho(a_1) = \dots = \rho(a_m)$ or $\rho(a_1) < \dots < \rho(a_m)$. If not, then we can find three consecutive candidates a_j , a_{j+1} , and a_{j+2} such that either $\rho(a_j) = \rho(a_{j+1}) < \rho(a_{j+2})$ or $\rho(a_j) < \rho(a_{j+1}) = \rho(a_{j+2})$. Both of these options are impossible in our case due to the existence of voters with preferences $a_{j+1} \succ a_j \succ a_{j+2}$ and $a_{j+1} \succ a_{j+2} \succ a_j$.

If $\rho(a_1) = \dots = \rho(a_m)$ the second condition is trivially true. Suppose $\rho(a_1) < \dots < \rho(a_m)$. Because voter 3 (resp. 4) prefers candidate a_{t-1} (resp. a_{t+1}) the most, we have $\rho(v_3) \in (\rho(a_{t-2}), \rho(a_t))$ (resp. $\rho(v_4) \in (\rho(a_t), \rho(a_{t+2}))$). We can now show

$$\begin{aligned}
d(a_1, a_m) &\leq d(a_1, v_4) + d(v_4, a_m) \\
&\leq 2d(a_1, v_4) \quad (\because a_m \succ_4 a_1) \\
&\leq 2d(a_1, a_{m-2}) \quad (\because \rho(a_t) < \rho(v_4) < \rho(a_{m-2})) \\
&\leq 2(d(a_1, v_3) + d(v_3, a_{m-2})) \\
&\leq 4d(a_1, v_3) \quad (\because a_{m-2} \succ_3 a_1) \\
&\leq 4d(a_1, a_t). \quad (\because \rho(a_1) < \rho(v_3) < \rho(a_t))
\end{aligned}$$

This concludes the proof. □

We can now restate our theorem:

For $m \geq 9$ and constant $\alpha \in (0, 0.5]$, $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}}^\alpha(\widehat{f}_{\text{fail}})$ is upper bounded by a constant, whereas $\phi_{\mathcal{I}_{m, \text{sep-}\mathbb{R}}}^\alpha(f_{\text{fail}})$ is unbounded.

Proof. First, we show that $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha}(f_{\text{fail}})$ is unbounded. Consider the following instance $I = (V, A, M, \rho, \pi)$. Let $V = \{v_1, \dots, v_{2n}\}$ (where $n \geq m^2$), $A = \{a_1, \dots, a_m\}$ (m being odd), and $M = (\mathbb{R}, d)$ with d being the Euclidean distance on the line.

The embedding function ρ is as follows. For $\ell \in [m]$, $\rho(a_\ell) = \frac{\ell-1}{m-1}$; that is, candidates a_1 through a_m are uniformly spaced in $[0, 1]$ with $\rho(a_1) = 0$ and $\rho(a_m) = 1$.

Fix $\epsilon < 1/m^2$. The voters are embedded as follows.

$$\begin{aligned} \rho(v_1) &= \rho(a_1) - \epsilon, \\ \rho(v_2) &= \rho(a_m) + \epsilon, \\ \rho(v_3) &= \rho(a_t - 1) + \epsilon, \\ \rho(v_4) &= \rho(a_t + 1) - \epsilon, \\ \rho(v_{4+(2j-1)}) &= \rho(a_{j+1}) + \epsilon && \forall j \in [m-2], \\ \rho(v_{4+2j}) &= \rho(a_{j+1}) - \epsilon && \forall j \in [m-2], \\ \rho(v_j) &= \frac{1}{2} && \forall j \geq 2m+1. \end{aligned}$$

Finally, in the party affiliation π , since we are just showing how bad the distortion of direct elections are and f is affiliation-independent, the party affiliation is not important, and the outcome is independent of π , so we can assign it in any way such that half the voters are of one party and half are of the other. This construction is all we need since $I \in \mathcal{I}_{m,\text{sep-}\mathbb{R}}^{0.5} \subseteq \mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha$ for all $\alpha \in (0, 0.5]$.

Next, it is also easy to check that election E^I falls under the special case of f_{fail} . Hence, $f_{\text{fail}}(E^I) = a_1$. Note that $C(a_1) \geq (2n - 2m) \cdot |0 - \frac{1}{2}| > n - m$ because a_1 is at distance $\frac{1}{2}$ from all but $2m$ voters located at $\frac{1}{2}$. In contrast, $C(a_t) \leq 2m \cdot 1$ because a_t is at zero distance from all but $2m$ voters (and its distance from those $2m$ voters is at most 1). Thus, $\phi(f_{\text{fail}}, I) \geq (n - m)/2m$. Since n is unbounded, $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha}(f_{\text{fail}})$ is also unbounded.

Finally, we show that $\phi_{\mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha}(\widehat{f}_{\text{fail}})$ is upper bounded by a constant. Fix an instance $I \in \mathcal{I}_{m,\text{sep-}\mathbb{R}}^\alpha$. For notational simplicity, we refer to the number of candidates in I as m , though the proof below works if it is less than m , too. First, assume $|\rho(A)| \neq 2$ (we will handle the case $|\rho(A)| = 2$ later). Without loss of generality, assume that $\rho(a_1) \leq \dots \leq \rho(a_m)$, and that for a fixed $q \in [m]$, candidates a_1, \dots, a_q are affiliated with party -1 and the rest are affiliated with party 1 .

Let I_{-1} and I_1 be the primary instances corresponding to I . Let a_{OPT} be an optimal candidate for I . Without loss of generality, suppose it is affiliated with party -1 .

In the proof of Theorem 2, $\phi(\widehat{f}, I)$ depends only on the distortion of f on the primary instance of the party that a_{OPT} is affiliated with. Hence, if primary election $E^{I_{-1}}$ does not fall under the special case of f_{fail} , then f_{fail} applies a Condorcet-consistent rule on I_{-1} , ensuring that $\phi(f, I_{-1})$ is at most 3. In this case, by Theorem 2, $\phi(\widehat{f}, I)$ is also upper bounded by a constant.

Suppose $E^{I_{-1}}$ falls under the special case of f_{fail} . Let $t = (q+1)/2$ and $d_{-1} = d(a_1, a_q)$. Then, by Lemma 9, $\min\{d(a_t, a_1), d(a_t, a_q)\} \geq d_{-1}/4$. From now on, we shall use asymptotic notation liberally for simplicity.

Recall that there is a set of voters $S \subset V_{-1}$ whose top candidate was a_t , and

$$|S| = |V_{-1}| - 2q = \Omega(|V_{-1}|) = \Omega(n),$$

where the second transition holds because in the special case, $|V_{-1}| \geq q^2$, and the final transition holds because $|V_{-1}| \geq \alpha n$.

Note that for every $i \in S$ and $j \in V_1$, $d(i, j) \geq d_{-1}/8$. And $|V_1| \geq \alpha n$. Hence, we have $\Omega(n)$ pairs of voters (i, j) such that $d(i, j) \geq d_{-1}/8$. Further, $d(a_{\text{OPT}}, i) + d(a_{\text{OPT}}, j) \geq$

$d(i, j)$. Hence, it follows that

$$C(a_{\text{OPT}}) = \Omega(n) \cdot d_{-1}. \quad (6)$$

Let $a^* = \widehat{f}(I)$. If $a^* = a_{-1}^* = a_1$, then we have

$$\begin{aligned} C(a^*) &\leq C(a_{\text{OPT}}) + n \cdot d(a^*, a_{\text{OPT}}) \\ &\leq C(a_{\text{OPT}}) + n \cdot d_{-1} = O(C(a_{\text{OPT}})), \end{aligned}$$

yielding a constant upper bound on $\phi(\widehat{f}_{\text{fail}}, I) = C(a^*)/C(a_{\text{OPT}})$.

On the other hand, if $a^* = a_1^*$, we have

$$\begin{aligned} C(a_{\text{OPT}}) &\geq C(a_1) - n \cdot d(a_1, a_{\text{OPT}}) \\ &\geq \frac{n}{2} \frac{d(a_1, a^*)}{2} - n \cdot d_{-1} \\ &\geq \frac{n}{2} \frac{d(a_{\text{OPT}}, a^*)}{2} - O(C(a_{\text{OPT}})). \end{aligned}$$

Here, the second transition follows because in the general election, at least $n/2$ voters vote for a^* over a_1 and $d(a_1, a_{\text{OPT}}) \leq d_{-1}$, and the final transition follows from Equation (6). This implies

$$C(a_{\text{OPT}}) = \Omega(n \cdot d(a_{\text{OPT}}, a^*)). \quad (7)$$

On the other hand, we have

$$C(a^*) \leq C(a_{\text{OPT}}) + n \cdot d(a_{\text{OPT}}, a^*) = O(C(a_{\text{OPT}})),$$

where the last transition follows due to Equation (7). Hence, we again have the desired constant upper bound on $\phi(\widehat{f}_{\text{fail}}, I)$.

Finally, if $\rho(A) = \{x_1, x_2\}$ (w.l.o.g., $x_1 < x_2$), due to separability, we have two options:

1. $\rho(A_{-1}) = x_1$ and $\rho(A_1) = x_2$: Therefore, $a_{\text{OPT}} \in \{a_{-1}^*, a_1^*\}$, and since a^* is the majority winner, $a^* = a_{\text{OPT}}$, and the distortion is 1.
2. $\rho(A_{-1}) = \{x_1, x_2\}$ and $\rho(A_1) = x_2$: If $a_{\text{OPT}} \in \{a_{-1}^*, a_1^*\}$, then it is similar to the previous case. Otherwise, this means $\rho(a_{\text{OPT}}) = x_1$ and $\rho(a_{-1}^*) = \rho(a_1^*) = x_2$. Separability means the voters of party 1 lie in $\{x | x \geq x_2\}$, so $C(a_{\text{OPT}}) \geq |V_1| \cdot d(x_1, x_2) \geq \alpha \cdot n \cdot d(x_1, x_2)$. While $C(a^*) \leq C(a_{\text{OPT}}) + |V_{-1}| \cdot d(x_1, x_2) \leq C(a_{\text{OPT}}) + n(1 - \alpha)d(x_1, x_2)$. Combining these two equations we get the distortion is $\frac{1+2\alpha}{\alpha}$.

□