

Every choice function is pro-con rationalizable

Serhat Dogan and Kemal Yildiz

Abstract

We consider an agent who is endowed with two sets of orderings: pro-orderings and con-orderings. For each choice set, if an alternative is the top-ranked by a *pro-ordering* (*con-ordering*), then this is a *pro* (*con*) for choosing that alternative. The alternative with more pros than cons is chosen from each choice set. Each ordering may have a *weight* reflecting its salience. In this case, each alternative is chosen with a probability proportional to the total weight of its pros and cons. We show that every nuance of the rich human choice behavior can be captured via this structured model. Our technique requires a generalization of Ford-Fulkerson Theorem, which may be of independent interest. As an application of our results, we show that every choice rule is plurality-rationalizable.

1 Introduction

Charles Darwin, the legendary naturalist, wrote “The day of days!” in his journal on November 11, 1838, when his cousin Emma Wedgwood accepted his marriage proposal. However, whether to marry at all had been a hard decision for Darwin. Just a few months prior, Darwin had scribbled a carefully considered list of *pros*—such as “constant companion” and “charms of music”—and *cons*—such as “fewer conversations with clever people” and “no books”—regarding the potential impact of marriage on his life.¹ With this list of pros and cons, Darwin seems to follow a choice procedure ascribed to Benjamin Franklin.² Here we present Franklin (1887)’s choice procedure in his own words.

To get over this, my Way is, to divide half a Sheet of Paper by a Line into two Columns, writing over the one Pro, and over the other Con. I endeavour to estimate their respective Weights; and where I find two, one on each side, that seem equal, I strike them both out: If I find a Reason pro equal to some two Reasons con, I strike out the three. If I judge some two Reasons con equal to some three Reasons pro, I strike out the five; and thus proceeding I find at length where the Ballance lies. And tho’ the Weight of Reasons cannot be taken with the Precision of Algebraic Quantities, yet when each is thus considered separately and comparatively, and the whole lies before me, I think I can judge better, and am less likely to take a rash Step; and in fact I have found great Advantage from this kind of Equation, in what may be called Moral or Prudential Algebra.

Choice models most commonly used in economics are based on maximization of preferences. An alternative mode of choice, which is common for the scholarly work in other social disciplines such as history, law, and political science, is the less formal *reason-based analysis* (Shafir et al. (1993)). Reason-based analysis is also commonly used for the analysis of ‘case studies’ in business and law schools.³ In the vein of Franklin’s prudential algebra, first, various arguments that support or oppose an alternative are identified, then the balance of these arguments determines the choice.⁴ We formulate and analyze the *pro-con choice model* that connects these two approaches by presenting a reason-based choice model, in which the ‘reasons’ are formed via a preference-based language.

¹See Glass (1988) for the full list.

²In 1772, Joseph Priestley wrote a letter to Benjamin Franklin asking for Franklin’s advice on a decision he was trying to make. Franklin wrote back indicating that he could not tell him what to do, but he could tell him how to make his decision, and suggested his *prudential algebra*.

³Köksalan et al. (2011) reports Franklin’s moral algebra as an early example of multi-criteria decision making.

⁴Shafir et al. (1993) argue that reason-based analyses have been used to understand unique historic, legal and political decisions. Examples include presidential decisions taken during the Cuban missile crisis (Allison (1971)), the Camp David accords (Telhami (1990)), and the Vietnam war (Gelb & Betts (2016)).

We formulate the pro-con choice model in the deterministic choice setup by extending Franklin’s prudential algebra to choice sets that possibly contain more than two alternatives. A (*deterministic*) *pro-con model* (pcM) is a pair $\langle \succ, \triangleright \rangle$ such that $\succ = \{\succ_1, \dots, \succ_m\}$ is a set of *pro-orderings* and $\triangleright = \{\triangleright_1, \dots, \triangleright_q\}$ is a set of *con-orderings*. We require that an ordering can not both serve as a pro- and con-ordering. Since \succ and \triangleright are defined as sets of orderings rather than lists or profiles of orderings, each ordering can be used only once as a pro- or con-ordering.⁵ Given an pcM $\langle \succ, \triangleright \rangle$, for each choice set S and alternative x , if x is the \succ_i -top-ranked alternative in S for some $\succ_i \in \succ$, then we interpret this as a ‘pro’ for choosing x from S . On the other hand, if x is the \triangleright_i -top-ranked alternative in S for some $\triangleright_i \in \triangleright$, then we interpret this as a ‘con’ for choosing x from S .

Our central new concept is the following: A *choice function*⁶ is *pro-con rational(izable)* if there is an pcM $\langle \succ, \triangleright \rangle$ such that for each choice set S , an alternative x is chosen from S if and only if pros for choosing x from S are more than the cons for choosing x from S .⁷

A *random pro-con model* (RpcM) is a triplet $\langle \succ, \triangleright, \lambda \rangle$, where \succ and \triangleright stand for the sets of pro-orderings and con-orderings, as before. The weight function λ assigns to each pro-ordering $\succ_i \in \succ$ a value in the $[0, 1]$ interval and con-ordering $\triangleright_i \in \triangleright$ a value in the $[-1, 0]$ interval, which we interpret as a measure of the salience of each ordering.⁸ The total weight of an alternative in a choice set is the total weight of the pro-and con-orderings at which it is top-ranked. To make a choice from each choice set, a pro-con-rational agent considers the alternatives with a positive total weight, and chooses each alternative from this consideration set with a probability proportional to its total weight.

The most familiar stochastic choice model in economics is the *random utility model* (RUM), which assumes that an agent is endowed with a probability measure μ over a set of orderings \succ such that he randomly selects an ordering to be maximized from \succ according to μ . An RUM $\langle \succ, \mu \rangle$ is an RpcM in which there is no set of con-orderings. Both the RpcM and the RUM are *additive* models, in the sense that the choice probability of an alternative is calculated by summing up the weights assigned to the orderings. The primitives of both the RpcM and RUM are *structurally invariant*, in the sense that the decision maker uses the same $\langle \succ, \mu \rangle$ and $\langle \succ, \triangleright, \lambda \rangle$ to make a choice from each choice set. These two features of RUM reflect themselves in its characterization.⁹ Despite the similarity between the RpcM and the RUM, in Theorem 2, we show that every random choice function is pro-con rational. Our technique is build on our Theorem 3 an original extension of Ford Jr & Fulkerson (2015)’s seminal result in optimization theory. Then, by using the construction in Theorem 2’s proof together with two key results from the integer-programming literature, in Theorem 1, we show that each (deterministic) choice function is pro-con rational.¹⁰

The remaining observations in the paper are as follows. In Section 2.3, we observe that our Theorem 1 fails to hold in the context of multi-valued choice rules unless we allow multiple appearance of an ordering as a pro- or con-ordering. In Section 2.4, we illustrate that our results facilitate identification of other inclusive choice models, by showing that each choice function is *plurality-rationalizable*. The model and the result can thought of as a generalization of an earlier model and a related result by McGarvey (1953). For the uniqueness of representation, the RpcM has characteristics similar to the RUM, which we present and discuss in Section 3.3.

⁵One concern is the number of orderings in \succ and \triangleright . It follows from this requirement that if there are n alternatives in X , then at most $n!$ -many orderings are used in a pro-con model.

⁶A *choice function* C singles out an alternative from each *choice set* S , which is a nonempty subset of the grand *alternative set* X .

⁷In extending Franklin’s prudential algebra, one can consider a sequential pro-con model in which first the alternatives that fail to have more pros than cons in the given choice set are eliminated, and then the elimination continues until an alternative is singled out. Our model is a specific sequential pro-con model in which all the alternatives but the chosen one are eliminated in the first step.

⁸In line with the experimental findings of Shafir (1993) indicating that the weight assigned to the pros is more than the weight assigned to the cons, we require the total weighted sum of pro-orderings and con-orderings be unity.

⁹Namely, the random choice functions that render a random utility representation are those with nonnegative Block-Marschak polynomials. See Block & Marschak (1960), Falmagne (1978), McFadden (1978), and Barberá & Pattanaik (1986).

¹⁰This result does not directly follow from Theorem 2, since a pro-con model is not a direct adaptation of the random pro-con model, in that we require each ordering to have a fixed unit weight instead of having fractional weights. To best of our knowledge the use of integer programming techniques in this context is new.

1.1 Related literature

In the deterministic choice literature, previous choice models proposed by [Kalai et al. \(2002\)](#) and [Bossert & Sprumont \(2013\)](#) yield similar “anything goes” results. A choice function is *rationalizable by multiple rationales* ([Kalai et al. \(2002\)](#)) if there is a collection of preference relations such that for each choice set the choice is made by maximizing one of these preferences. Put differently, the decision maker selects an ordering to be maximized for each choice set. A choice function is *backwards-induction rationalizable* ([Bossert & Sprumont \(2013\)](#)) if there is an extensive-form game such that for each choice set the backwards-induction outcome of the restriction of the game to the choice set coincides with the choice. In this model, for each choice set, a new game is obtained by pruning the original tree of all branches leading to unavailable alternatives. For random choice functions, [Manzini & Mariotti \(2014\)](#) provide an anything-goes result for the *menu-dependent random consideration set rules*, in which an agent keeps a single preference relation and attaches to each alternative a choice-set-specific attention parameter. Then, he chooses an alternative with the probability that no more-preferable alternative grabs his attention. In contrast to these models, we believe that the pro-con model is more structured and exhibits limited context dependency. An agent following a pro-con model restricts the pro- and con-orderings to the given choice set to make a choice.

It may be of interest to view our model from the perspective of probabilistic social choice. Existing work in this literature show that the class of probabilistic group decision rules have considerable richness and appeal. As a partial list one can consider [Intriligator \(1973\)](#), [Barberá & Sonnenschein \(1978\)](#), [Pattanaik & Peleg \(1986\)](#), and [Intriligator \(1982\)](#). These studies typically investigate the structure of coalitional power under probabilistic social decision rules. The closest to our work is [Pattanaik & Peleg \(1986\)](#) who axiomatically characterize the random dictatorship procedure, in which there is a probability measure μ on the members of the society N such that for each profile of individual preferences $\{\succ_i\}_{i \in N}$, the society chooses from each choice set according to the RUM $\langle \{\succ_i\}_{i \in N}, \mu \rangle$. Along these lines, for a social-choice interpretation of the mixed-sign representation in an RpcM, consider a chair who stochastically aggregates different opinions in a committee to make a choice. It is typically assumed that as more committee members top rank an alternative, the choice probability of this alternative increases. However, there may be an antagonistic relationship between the chair and some committee members, so that the chair would be less likely to choose the alternative favored by them.

Franklin’s moral algebra is well known in the field of multi-criteria decision making.¹¹ [Köksalan et al. \(2011\)](#) reports Franklin’s moral algebra as one of the earliest examples of multi-criteria decision making. [Hammond et al. \(1998\)](#) proposes the related ‘even swaps’ method as a practical way of making trade-offs among any set of objectives across a range of alternatives. Our approach diverges from the bulk of research in this literature in taking the route of revealed preference analysis. That is, we recover the primitive objectives and the way that they are reconciled from observed choices.

Our Theorem 2 is related to a result in a contemporary paper by [Saito \(2017\)](#), who offers characterizations of the mixed logit model. It follows from the results of this paper, which is proved by using a different approach, that each RCF can be expressed as a convex combination of two random utility functions.¹² We discuss the technical differences at the end of Section 3.2.

2 Deterministic pro-con choice

2.1 The model

Given a nonempty finite alternative set X , any nonempty subset S is called a **choice set**. Let Ω denote the collection of all choice sets. A (deterministic) choice function C is a mapping that assigns each choice set $S \in \Omega$ a member of S , that is $C : \Omega \rightarrow X$ such that $C(S) \in S$. An **ordering**, denoted generically by \succ_i or \triangleright_i , is a complete, transitive, and antisymmetric binary relation on X .

¹¹For a classical reference one can consult [Arrow et al. \(1986\)](#).

¹²We are grateful to an anonymous referee for bringing this connection to our awareness.

A **(deterministic) pro-con model (pcM)** is a pair $\langle \succ, \triangleright \rangle$, where $\succ = \{\succ_1, \dots, \succ_m\}$ and $\triangleright = \{\triangleright_1, \dots, \triangleright_q\}$ are sets of pro- and con-orderings on X . Given an pcM $\langle \succ, \triangleright \rangle$, for each choice set S and alternative $x \in S$, if x is the \succ_i -top-ranked alternative in S for some $\succ_i \in \succ$, then we interpret this as a ‘pro’ for choosing x from S . On the other hand, if x is the \triangleright_i -top-ranked alternative in S for some $\triangleright_i \in \triangleright$, then we interpret this as a ‘con’ for choosing x from S . Define $Pros(x, S) = \{\succ_i \in \succ : x = \max(S, \succ_i)\}$ and $Cons(x, S) = \{\triangleright_i \in \triangleright : x = \max(S, \triangleright_i)\}$.

Definition 1 A choice function C is **pro-con rational** if there is an pcM $\langle \succ, \triangleright \rangle$ such that for each choice set $S \in \Omega$ and $x \in S$, $C(S) = x$ if and only if $|Pros(x, S)| > |Cons(x, S)|$.

Note that if an agent is pro-con rational, then at each choice set S there should be a single alternative x such that the number of $Pros(x, S)$ is greater than the number of $Cons(x, S)$. Moreover, the pro-con model is not a direct adaptation of its random counterpart. In that, we require each ordering to have a fixed unit weight, instead of having fractional weights. Next, to illustrate how the model works, we revisit Luce and Raiffa’s dinner example (Luce & Raiffa (1957)) by following a pro-con model.

Example 1 Suppose you choose chicken when the menu consists of steak and chicken only, yet go for the steak when the menu consists of steak (S), chicken (C), and fish (F). Consider the pro-orderings \succ_1 and \succ_2 that order the three dishes according to their *attractiveness* and *healthiness*, so suppose $S \succ_1 F \succ_1 C$ and $F \succ_2 C \succ_2 S$. As a con-ordering, consider $F \triangleright S \triangleright C$, which orders the dishes according to their *riskiness*. Since cooking fish requires expertise, it is the most risky one and chicken is the safest option.

Now, to make a choice from the grand menu, the pros are: “ S is the most attractive”, “ F is the most healthy”, but also “ F is the most risky”. Thus, S is chosen from the grand menu. If only S and C are available, then we have “ C is the most healthy”, “ S is the most attractive”, but also “ S is the most risky”, so C is chosen.

In our Definition 1, we ask for a rather structured representation that corresponds to one-to-one elimination in Franklin’s prudential algebra. We see at least two benefits of this stringency. First, we obtain the uniqueness property presented in Section 3.3. Second, in Section 2.3, we argue that our Theorem 1 fails to hold in the context of multi-valued choice rules. Finally, given our Theorem 1, one can use our representation to identify other inclusive choice models, which otherwise may not be an easy exercise. In Section 2.4, we present an application along these lines, in which we show that each choice function is plurality-rationalizable.

2.2 Main result

We show that every choice function is pro-con rational. In the language of mathematical programming, in Theorem 2, we show that the relaxed (convex) problem has a solution. In Section 5.3, we prove Theorem 1, which translates into finding an integer solution, by using the construction in Theorem 2’s proof together with two key results from the integer-programming literature, the ones developed by Hoffman & Kruskal (2010) and Heller & Tompkins (1956).

Theorem 1 Every choice function is pro-con rational.

Remark 1 The constructed pro-con representation is a rather parsimonious one. To see this, consider a more stringent pro-con model, in which if an alternative x is chosen from a choice set S , it is barely chosen in the sense $|Pros(x, S)| - |Cons(x, S)| = 1$, and if an alternative y is not chosen, it is barely not chosen in the sense $|Pros(y, S)| - |Cons(y, S)| = 0$. It follows from the proof of Theorem 1 that the same anything-goes-result holds for this model.

2.3 Extension to multi-valued choice

There are instances in which an agent must choose more than a single alternative from a choice set. For example, consider a school that chooses a cohort from a set of applicants or a professor who chooses a set of questions out of his archive to prepare an exam. As for the random choice, imagine that we have access the support of the random choice function, but not the frequencies, then the observed choice behavior yields a choice rule.¹³

So far, we have assumed that the observed choice behavior is summarized by a choice function or a random choice rule. Both models rule out the possibility that choice can be multi-valued. Formally, a **choice rule** $\mathbb{C} : \Omega \rightarrow \Omega$ such that for each $S \in \Omega$, $\mathbb{C}(S) \subset S$. A choice rule is pro-con rational if there exists a pro-con model $\langle \succ, \triangleright \rangle$ such that for each choice set $S \in \Omega$, $\mathbb{C}(S) = \arg \max_{x \in S} (|Pros(x, S)| - |Cons(x, S)|)$. That is, for each choice set $S \in \Omega$ and $x \in S$, $x \in \mathbb{C}(S)$ if and only if $|Pros(x, S)| - |Cons(x, S)| \geq |Pros(y, S)| - |Cons(y, S)|$ for each $y \in S$.

A natural question is if our result in Theorem 1 extends to choice rules or not. To see that not every choice rule is pro-con rational, consider the choice rule \mathbb{C} defined on $\{x, y, z\}$ such that $\mathbb{C}(\{x, y, z\}) = \{x, y\}$, $\mathbb{C}(\{x, y\}) = \{x\}$, $\mathbb{C}(\{y, z\}) = \{y\}$, and $\mathbb{C}(\{x, z\}) = \{z\}$. It is easy to see that \mathbb{C} is not pro-con rational.¹⁴ The stringency in here derives from the requirement that each ordering can be used only once as a pro- or con-ordering in a pro-con model.

In contrast, if we allow multiple appearance of an ordering as a pro- or con-ordering, then every choice rule can be recovered. To see this, let \mathbb{C} be a choice rule, and let p be the associated random choice function such that for each $S \in \Omega$ and $x \in \mathbb{C}(S)$, $p(x, S) = 1/|\mathbb{C}(S)|$. It follows from Theorem 2 that there is a random pro-con model $\langle \succ, \triangleright, \lambda \rangle$ which represents p . Moreover, it follows from the construction in the proof of Theorem 2 that if for each $S \in \Omega$ and $x \in \mathbb{C}(S)$, $p(x, S)$ is a rational number, then for each $\succ_i \in \succ$ and $\triangleright_j \in \triangleright$, we can choose $\lambda(\succ_i) = m_i/M$ and $\lambda(\triangleright_j) = m_j/M$, where m_i, m_j, M are positive integers. Now, consider a list (or a profile) of pro-orderings with m_i -many copies of \succ_i and m_j -many copies of \triangleright_j for each $\succ_i \in \succ$ and $\triangleright_j \in \triangleright$. It directly follows from this construction that for each $S \in \Omega$ and $x \in S$, $x \in \mathbb{C}(S)$ if and only if x maximizes the difference between number of pro-orderings at which x is top-ranked in S and the number of con-orderings at which x is top-ranked in S .

2.4 Plurality-rationalizable choice rules

We analyze a collective decision making model based on plurality voting. It turns out that this model is closely related to our pro-con choice model. To introduce this model, let $[\succ^*] = [\succ_1^*, \dots, \succ_m^*]$ be a preference profile, which is a list of orderings. In contrast to a set of orderings, denoted by \succ or \triangleright , an ordering \succ_i^* can appear more than once in a preference profile $[\succ^*]$. For each choice set $S \in \Omega$ and $x \in S$, x is a **plurality winner of $[\succ^*]$ in S** if for each $y \in S \setminus \{x\}$, the number of orderings in $[\succ^*]$ that top rank x in S is more than or equal to the number of orderings in $[\succ^*]$ that top rank y in S . That is, for each $y \in S \setminus \{x\}$, $|\{\succ_i^* \in [\succ^*] : x = \max(S, \succ_i^*)\}| \geq |\{\succ_i^* \in [\succ^*] : y = \max(S, \succ_i^*)\}|$. Next, we define plurality-rationalizability, then by using our Theorem 1, we show that every choice rule is plurality-rationalizable.

Definition 2 A choice rule C is **plurality-rationalizable** if there is preference profile $[\succ^*]$ such that for each choice set $S \in \Omega$ and $x \in S$, $x \in \mathbb{C}(S)$ if and only if x is a plurality winner of $[\succ^*]$ in S .

Proposition 1 Every choice rule is plurality-rationalizable.¹⁵

¹³See, for example, Fishburn (1978) who explores a connection in this vein.

¹⁴The two stage threshold representation analyzed by Manzini et al. (2013) has a similar feature. In that, although each choice function has a two-stage threshold representation, this does not hold for choice rules. That is, for each choice function there is a triplet $\langle f, \theta, g \rangle$ such that for each $S \in \Omega$, the alternative that maximizes $g(x)$ subject to $f(x) \geq \theta(S)$ is chosen. However, such a two stage threshold representation can not be obtained for every choice rule.

¹⁵Our initial result was for choice functions. We thank Vicki Knoblauch and an anonymous referee for suggesting the extension to choice rules.

Proof. Let \mathbb{C} be a choice rule. In Section 2.3, by using Theorem 1, we show that if we allow multiple appearance of an ordering as a pro- or con-ordering, then every choice rule is pro-con rational. First, to formalize this representation, let \succ and \triangleright be the set of pro- and con-orderings such that each $\succ_i \in \succ$ ($\triangleright_i \in \triangleright$) is copied k_i times to represent \mathbb{C} . Then, define for each $S \in \Omega$ and $x \in S$, $SPros(x, S) = \sum_{\{\succ_i \in Pros(x, S)\}} k_i$ and $SCons(x, S) = \sum_{\{\triangleright_i \in Cons(x, S)\}} k_i$, where $Pros(x, S)$ and $Cons(x, S)$ are defined as usual with respect to \succ and \triangleright . Now, we know that for each $S \in \Omega$ and $x \in S$, $x \in \mathbb{C}(S)$ if and only if $x \in \arg \max_{x \in S} (|SPros(x, S, \succ^*)| - |SCons(x, S, \triangleright^*)|)$.

Now, to construct the desired preference profile, let $k = \max_{\{\triangleright_i \in \triangleright^*\}} k_i$, and begin with the list of all orderings defined on X copied k times. This is preference profile with $kn!$ elements. Then, eliminate k_i copies of the inverse of each ordering $\triangleright_i \in \triangleright$, and add k_i copies of each ordering $\succ_i \in \succ$. Note that since we have k copies of each ordering, the elimination part is well-defined. Let $[\succ^*]$ be the obtained preference profile.

We show that for each $S \in \Omega$ and $x \in S$, $x \in \mathbb{C}(S)$ if and only if x is a plurality winner of $[\succ^*]$ in S . We know that $x \in \mathbb{C}(S)$ if and only if for each $y \in S \setminus \{x\}$, $|SPros(x, S)| - |SCons(x, S)| \geq |SPros(y, S)| - |SCons(y, S)|$. Now, note that by construction of $[\succ^*]$, for each $y \in S$ the number of orderings in $[\succ^*]$ that top rank y in S equals k times the number of all orderings that top rank y in S , added to $|SPros(y, S)| - |SCons(y, S)|$. Since for each $y \in S$, the number of all orderings that top rank y in S is fixed, it follows that $x \in \mathbb{C}(S)$ if and only if x is a plurality winner of $[\succ^*]$ in S . ■

If we restrict our attention to choice functions, then we can consider an even more stringent model. In which, we require that an alternative x is chosen from a choice set S if and only if x is the plurality winner *at the margin*, in the sense that if x receives k votes then each other alternative receives $k-1$ votes. It follows from Remark 1 and the proof of Proposition 1 that every choice function is plurality-rationalizable via this more demanding model.

In an early paper McGarvey (1953) shows that for each asymmetric and complete binary relation, there exists a preference profile such that the given binary relation is obtained from the preference profile by comparing each pair of alternatives via majority voting.¹⁶ We obtain McGarvey's result, as a corollary to Proposition 1. To see this, note that if we restrict a choice rule to binary choice sets, then we obtain an asymmetric and complete binary relation. Since for binary choices, being a plurality winner means being a majority winner, McGarvey's result directly follows.

3 Random pro-con choice

3.1 The model

A **random choice function** (RCF) p is a mapping that assigns each choice set $S \in \Omega$, a probability measure over S . For each $S \in \Omega$ and $x \in S$, we denote by $p(x, S)$ the probability that alternative x is chosen from choice set S .

A **random pro-con model** (RpcM) is a triplet $\langle \succ, \triangleright, \lambda \rangle$, where \succ and \triangleright stand for the sets of pro- and con-orderings on X as before. The **weight function**, denoted by λ , is such that for each $\succ_i \in \succ$ and $\triangleright_i \in \triangleright$, we have $\lambda(\succ_i) \in (0, 1]$, $\lambda(\triangleright_i) \in [-1, 0)$, and the weighted sum of pro-orderings and con-orderings is one, i.e. $\sum_{\{\succ_i \in \succ\}} \lambda(\succ_i) + \sum_{\{\triangleright_i \in \triangleright\}} \lambda(\triangleright_i) = 1$. The weight function λ acts like a probability measure over the set of orderings that can assign negative values. In measure theoretic language, the primitive of a random pro-con model is a *signed probability measure* defined over the set of orderings.

Given an RpcM $\langle \succ, \triangleright, \lambda \rangle$, for each choice set S and alternative $x \in S$, if x is the \succ_i -top-ranked alternative in S for some $\succ_i \in \succ$, then we interpret this as a 'pro' for choosing x from S . On the other hand, if x is the \triangleright_i -top-ranked alternative in S for some $\triangleright_i \in \triangleright$, then we interpret this as a 'con' for choosing x from S . We interpret the weight assigned to each pro-ordering or con-ordering as a measure of the strength of that ordering.

¹⁶Stearns (1959) finds upper and lower bounds on the number of voters to generate any binary relation. Knoblauch (2016) provides an extension for infinite sets.

To define when an RCF is pro-con rational, let $Pros(x, S) = \{\succ_i \in \succ : x = \max(S, \succ_i)\}$ and $Cons(x, S) = \{\triangleright_i \in \triangleright : x = \max(S, \triangleright_i)\}$. Next, we formally define when an RCF is pro-con rational. For a given RpcM $\langle \succ, \triangleright, \lambda \rangle$, for each choice set $S \in \Omega$ and $x \in S$, we denote the total weight of x in S by $\lambda(x, S)$, i.e. $\lambda(x, S) = \lambda(Pros(x, S)) + \lambda(Cons(x, S))$. For each choice set $S \in \Omega$, let S^+ be the set of alternatives in S that receives a positive total weight, i.e. $S^+ = \{x \in S : \lambda(x, S) > 0\}$.

Definition 3 An RCF p is **pro-con rational** if there is an RpcM $\langle \succ, \triangleright, \lambda \rangle$ such that for each choice set $S \in \Omega$ and $x \in S$,

$$p(x, S) = \max \left\{ 0, \frac{\lambda(x, S)}{\sum_{\{y \in S^+\}} \lambda(y, S)} \right\} \quad (1)$$

That is, to make a choice from each choice set S , a pro-con-rational agent considers the alternatives with a positive total weight, and chooses each alternative from this consideration set with a probability proportional to its total weight. An equivalent formulation is as follows. An RCF p is pro-con rational if there is an RpcM $\langle \succ, \triangleright, \lambda \rangle$ such that for each choice set $S \in \Omega$ and $x \in S$, $p(x, S) = \lambda(Pros(x, S)) + \lambda(Cons(x, S))$, where $\lambda(Pros(x, S))$ and $\lambda(Cons(x, S))$ are the sum of the weights over $Pros(x, S)$ and $Cons(x, S)$. Our proof of Theorem 2 clarifies this equivalence.

As an alternative to the *random utility model* (RUM), [Tversky \(1972\)](#) proposes *elimination by aspects* model in which an agent views each alternative as a set of attributes. Then, at each stage, the agent selects an attribute with probability proportional to its weight and eliminates all the alternatives without the selected attribute. Pro-con model offers a choice procedure that both carries the act of selecting an ordering to be maximized as in the random utility model and elimination of the alternatives based on their attributes as in [Tversky \(1972\)](#)'s *elimination by aspects*. In that, a pro-con-rational agent's attitude to the relevant attributes is twofold: If it is a pro-ordering, then he seeks maximization as in the RUM, if it is a con-ordering, then he is satisfied by elimination of the worst alternative as in the *elimination by aspects* model. In line with this interpretation, we illustrate in [Example 2](#) that each ordering in an RpcM can be interpreted as an attribute or a relevant criterion.

To illustrate how RpcM works, we focus on a particular choice problem in which there are only two orderings (\succ_1, \succ_2) that are relevant for choice, such as price and quality, and present an *attraction effect* scenario.¹⁷ In this scenario, when we introduce an asymmetrically dominated alternative, called a *decoy*, the choice probability of the dominating alternative goes up. This choice behavior, known as the *attraction effect*, is incompatible with any RUM.

Example 2 (Attraction Effect) Suppose $X = \{x, y, z\}$, where x and y are two competing alternatives such that none clearly dominates the other, and z is another alternative that is dominated by x but not y . Consider the following RpcM $\langle \succ, \triangleright, \lambda \rangle$, in which there is single pair of orderings used both as the pro- and con-orderings, with weights shown in parenthesis. We can interpret this ordering pair as two distinct criteria that order the alternatives.

(1)	(1)	$(-\frac{1}{2})$	$(-\frac{1}{2})$
\succ_1	\succ_2	\succ_1	\succ_2
x	y	y	z
z	x	z	x
y	z	x	y

Now, since for both criteria x is better than z , we get $p(x, \{x, z\}) = 1$. Since x and y fail to dominate each other, and y fail to dominate z , we get $p(y, \{x, y\}) = p(y, \{y, z\}) = 1/2$. That is, z is a 'decoy' for x when y is available. Note that when only x and y are available, since x is the \succ_2 -worst alternative, x is eliminated with a weight of $1/2$. However, when the decoy z is added to the choice set, then x is no longer the \succ_2 -worst alternative, and we get $p(x, \{x, y, z\}) = 2/3$. That is, availability of decoy z increases the choice probability of x . Thus, our model captures the intuition that the choice probability

¹⁷Experimental evidence for the attraction effect is first presented by [Payne & Puto \(1982\)](#) and [Huber & Puto \(1983\)](#). Following their work, evidence for the attraction effect has been observed in a wide variety of settings. For a list of these results, consult [Rieskamp et al. \(2006\)](#).

of an alternative may increase when a *decoy* is added, since this alternative may no longer be the worst one according to a relevant attribute.

3.2 Main result

In our main result, we show that every random choice function is pro-con rational. We present a detailed discussion of the result in the introduction. We present the proof in Section 5.2. As a notable technical contribution, we generalize the Ford-Fulkerson Theorem (Ford Jr & Fulkerson (2015)) from combinatorial matrix theory to prove the result. Next, we state the theorem and present an overview of the proof. Then, we discuss the technical connection to Saito (2017).

Theorem 2 *Every random choice function is pro-con rational.*

An overview of the proof: For a given RCF p , we show that there is a *signed weight function* λ , which assigns each ordering \succ_i , a value $\lambda(\succ_i) \in [-1, 1]$ such that λ represents p . That is, for each choice set S and $x \in S$, $p(x, S)$ is the sum of the weights over orderings at which x is the top-ranked alternative. We prove this by induction.

To clarify the induction argument, for $k = 1$, let $\Omega_1 = \{X\}$ and let \mathcal{P}^1 consists of n -many equivalence classes such that each class contains all the orderings that top rank the same alternative, irrespective of whether these are chosen with positive probability. That is, for $X = \{x_1, \dots, x_n\}$, we have $\mathcal{P}^1 = \{[\succ^{x_1}], \dots, [\succ^{x_n}]\}$, where for each $i \in \{1, \dots, n\}$ and ordering $\succ_i \in [\succ^{x_i}]$, $x_i = \max(X, \succ_i)$. Now for each $x_i \in X$, define $\lambda^1([\succ^{x_i}]) = p(x_i, X)$. It directly follows that λ^1 is a signed weight function over \mathcal{P}^1 that represents the restriction of the given RCF to Ω_1 , denoted by p_1 . By proceeding inductively, it remains to show that we can construct λ^{k+1} over \mathcal{P}^{k+1} that represents p_{k+1} .

In Step 1 of the proof we show that finding such a λ^{k+1} boils down to finding a solution to the system of equalities described by *row sums* (RS) and *column sums* (CS). Up to this point the proof structure is similar to the one followed by Falmagne (1978) and Barberá & Pattanaik (1986) for the characterization of RUM.

To understand (RS), while moving from the k^{th} -step to the $(k + 1)^{th}$ -step, each $[\succ^k]$ is decomposed into a collection $\{[\succ_j^{k+1}]\}_{j \in J}$ such that for each $[\succ_j^{k+1}]$ there exists an alternative x_j that is not linearly ordered by $[\succ^k]$, but placed at $[\succ_j^{k+1}]$ right on top of the alternatives that are not linearly ordered by $[\succ^k]$. Therefore, the sum of the weights assigned to $\{[\succ_j^{k+1}]\}_{j \in J}$ should be equal to the weight assigned to $[\succ^k]$. This gives us the set of equalities formulated in (RS). To understand (CS), let S be the set of alternatives that are not linearly ordered by $[\succ^k]$. Now, we should design λ^{k+1} such that for each $x_j \in S$, $p(x_j, S)$ should be equal to the sum of the weights assigned to orderings at which x_j is the top-ranked alternative in S . The set of equalities formulated in (CS) guarantees this. This follows from our Lemma 2, which we obtain by using the *Mobius inversion*.¹⁸

Our proof is based on two interwoven observations. To understand the first, let us turn back to the induction argument. It is easy to see that the signed weight function λ^2 over \mathcal{P}^2 that represents p_2 is determined uniquely. That is, there is a unique λ^2 that satisfies equalities (RS) and (CS) formed for $k = 2$. But, then for λ^3 (in general for each $k \geq 3$) to be defined over \mathcal{P}^3 , the solution to the associated (RS) and (CS) for $k = 3$ is no longer unique. The difficulty is that although any λ^3 that satisfies equalities (RS) and (CS) for the $k = 3$ represents p_3 , depending on the choice of λ^3 , the (RS) and (CS) formed for a future step, $k > 3$, may not have a solution. Therefore, to conclude the induction successfully, for each $k \geq 3$, we should be “forwarding looking” in choosing λ^k .

Our second critical observation is that finding a solution to the system described by (RS) and (CS) can be translated to the following basic problem: Let $R = [r_1, \dots, r_m]$ and $C = [c_1, \dots, c_n]$ be two real-valued vectors such that the sum of R equals to the sum of C . Now, for which R and C can we find an $m \times n$ matrix $A = [a_{ij}]$ such that A has row sum vector R and column sum vector C , with each entry $a_{ij} \in [-1, 1]$? Ford Jr & Fulkerson (2015) provide a full answer to this question when R and C are positive real valued.¹⁹ However, there are two issues peculiar to our problem. First issue is

¹⁸Fiorini (2004) is the first who makes the same observation.

¹⁹Brualdi & Ryser (1991) provides a detailed account of similar results.

that the row and column sums can be negative real valued. Indeed, we get nonnegative-valued rows and columns only if the Block-Marschak polynomials are nonnegative, that is, the given p is an RUM. Second issue is that, related to our previous observation, we need “forward looking” solutions. In our Theorem 3, we provide a generalization of Ford-Fulkerson Theorem that paves the way for our proof by solving the two issues.

Theorem 3 (Generalized Ford-Fulkerson Theorem) *Let $R = [r_1, \dots, r_m]$ and $C = [c_1, \dots, c_n]$ be real-valued vectors with $-1 \leq r_i \leq 1$ and $-m \leq c_j \leq m$ such that $\sum_{i=1}^m r_i = \sum_{j=1}^n c_j$. If $2m \geq \sum_{i=1}^m |r_i| + \sum_{j=1}^n |c_j|$, then there is an $m \times n$ matrix $A = [a_{ij}]$ such that:*

- i. A has row sum vector R and column sum vector C ,
- ii. each entry $a_{ij} \in [-1, 1]$, and
- iii. for each $j \in \{1, \dots, n\}$, $\sum_{i=1}^m |a_{ij}| \leq |c_j| + \max\{0, \frac{\sum_{i=1}^m |r_i| - \sum_{j=1}^n |c_j|}{n}\}$.

To get an intuition for Theorem 3,²⁰ it is easy to see that the sum of the absolute values of the rows and columns should be bounded in order to extend the result to real-valued vectors. So, in Theorem 3, we require this sum be less than or equal to $2m$, where m is the number of rows. The choice of this specific bound has two implications. First, we can extend Ford-Fulkerson Theorem with real-valued rows and columns. This solves the first issue. Second, we guarantee that there is a solution that satisfy the bound in item (iii) of Theorem 3. This solution turns out to be the forwarding looking solution, which solves the second issue.

The rest of the proof is as follows. In Step 2, we show that (RS) equals (CS). In Step 3, by using a structural result presented in Lemma 3, we show that the row and column vectors associated with (RS) and (CS) satisfy the premises of our Theorem 3. This completes the construction of the desired signed weight function.

As discussed in Section 1.1, Saito (2017) independently shows that each RCF can be expressed as a convex combination of two random utility functions by using different techniques. To see our technical contribution note that by following the construction in our proof and directly applying the Ford-Fulkerson Theorem, each RCF can be expressed as an affine combination of random utility functions. To show that these weights can be chosen from $[-1, 1]$ interval, we generalize the Ford-Fulkerson Theorem (see Theorem 3) and follow a deliberate induction argument supported by other structural results, such as Lemma 3. We believe that our technique can be fruitful in solving similar random choice problems.

3.3 Uniqueness

An RCF may have different random utility representations even with disjoint sets of orderings. Falmagne (1978) argues that random utility representation is essentially unique. That is, the sum of the probabilities assigned to the orderings at which an alternative x is the k^{th} -top-ranked in a choice set is the same for all random utility representations of the given RCF. Similarly, the primitives of an RpcM are structurally invariant in the sense that the agent uses the same triplet $\langle \succ, \triangleright, \lambda \rangle$ to make a choice from each choice set. As an instance of this similarity, both models render a unique representation when there are only three alternatives.²¹ As for the general case, Proposition 2 provides a uniqueness result for the RpcM, which can be thought as the counterpart of Falmagne’s result for the RUM. Finally, as a direct corollary to Proposition 2, we present the counterpart of our uniqueness result for a pro-con model.

For a given RpcM $\langle \succ, \triangleright, \lambda \rangle$, let for each $S \in \Omega$ and $x \in S$, $\lambda(x = B_k | S, \succ, \triangleright)$ be the sum of the weights assigned to the pro- and con-orderings at which x is the k^{th} -top-ranked alternative in S . In our next result, we show that for each RCF the sum of the weights assigned to the orderings at which x is the k^{th} -top-ranked alternative in S is the same for each pro-con representation of the given RCF. That is, $\lambda(x = B_k | S, \succ, \triangleright)$ is fixed for each RpcM $\langle \succ, \triangleright, \lambda \rangle$ that represents the given RCF.

²⁰We present the proof in Section 5.1.

²¹This directly follows from the construction used to establish the base of induction in Theorem 2’s proof.

Proposition 2 *If $\langle \succ, \triangleright, \lambda \rangle$ and $\langle \succ', \triangleright', \lambda' \rangle$ are random pro-con representations of the same RCF p , then for each $S \in \Omega$ and $x \in S$,*

$$\lambda(x = B_k | S, \succ, \triangleright) = \lambda'(x = B_k | S, \succ', \triangleright'). \quad (2)$$

Proof. Let $\langle \succ, \triangleright, \lambda \rangle$ and $\langle \succ', \triangleright', \lambda' \rangle$ be two RPMs that represent the same RCF p . Now, for each choice set $S \in \Omega$, both λ and λ' should satisfy the identity (CS) used in Step 1 of the proof of Theorem 2. That is, for each $S \in \Omega$ and $x \in S$ both λ and λ' generates the same $q(x, S)$ value. Therefore, if we can show that $\lambda(x = B_k | S, \succ, \triangleright)$ can be expressed in terms of $q(x, \cdot)$, then (2) follows. To see this, let $\langle \succ, \triangleright, \lambda \rangle$ be any Rpm that represents p . Next, for each $S \in \Omega$, $x \in S$, and $k \in \{1, \dots, |S|\}$, consider a partition (S_1, S_2) of S such that $x \in S_2$ and $|S_1| = k - 1$. Let $\mathbb{P}(S, x, k)$ be the collection of all these partitions. Now, for each fixed $(S_1, S_2) \in \mathbb{P}(S, x, k)$, let $\lambda(x | S_1, S_2, \succ, \triangleright)$ be the sum of the weights of the orderings at which x is the top-ranked alternative in S_2 and the top-ranked alternative in S_1 . Note that for each such ordering, x is the k^{th} -top-ranked alternative in S . Now, it follows that we have:

$$\lambda(x = B_k | S, \succ, \triangleright) = \sum_{\{(S_1, S_2) \in \mathbb{P}(S, x, k)\}} \lambda(x | S_1, S_2, \succ, \triangleright). \quad (3)$$

Since for each $T \in \Omega$ such that $S_2 \subset T$ and $T \subset X \setminus S_1$, by definition, $q(x, T)$ gives the total weight of the orderings at which x is the top-ranked alternative in S , it follows that

$$\sum_{\mathbb{P}(S, x, k)} \lambda(x | S_1, S_2, \succ, \triangleright) = \sum_{\mathbb{P}(S, x, k)} \sum_{S_2 \subset T \subset X \setminus S_1} q(x, T). \quad (4)$$

Finally, if we substitute (3) in (4), then we express $\lambda(x = B_k | S, \succ, \triangleright)$ only in terms of $q(x, \cdot)$, as desired. ■

Next, we present the counterpart of our uniqueness result for a pro-con model. For a given Rpm $\langle \succ, \triangleright \rangle$, let for each $S \in \Omega$ and $x \in S$, $Pros^k(x, S)$ be the set of pro-orderings at which x is the k^{th} -top-ranked alternative in S . Similarly, let $Cons^k(x, S)$ be the set of con-orderings at which x is the k^{th} -top-ranked alternative in S . We show that for a given choice function, the difference between the number of pro-orderings at which x is the k^{th} -top-ranked alternative in S and the number of con-orderings at which x is the k^{th} -top-ranked alternative in S is the same for each pro-con representation of the given choice function. We obtain this result as a direct corollary to our Proposition 2.

Corollary 1 *If $\langle \succ, \triangleright \rangle$ and $\langle \succ', \triangleright' \rangle$ are pro-con representations of the same choice function C , then for each $S \in \Omega$, $x \in S$, and $k \in \{1, \dots, n\}$, both representations lead the same $|Pros^k(x, S)| - |Cons^k(x, S)|$ value.*

Proof. Since each pro- and con-ordering has a unit weight at each pro-con representation of a given choice function, $|Pros^k(x, S)| - |Cons^k(x, S)|$ equals $\lambda(x = B_k | S, \succ, \triangleright)$. Then, it follows from Proposition 2 that $|Pros^k(x, S)| - |Cons^k(x, S)|$ is fixed for each pro-con representation. ■

4 Conclusion

Our main results show that the pro-con model—an additive model similar to the RUM—provides a language to describe any choice behavior in terms of structurally-invariant primitives. The structural invariance of the pro-con model reflects itself as a form of uniqueness, which is similar to the uniqueness of a random utility model. Knowing that each choice function is pro-con rational facilitates identification of other permissive choice models. We present an application along these lines, in which we show that each choice rule is plurality-rationalizable. Although our study covers a rather extensive treatment of the pro-con model, we can hardly claim that it is exhaustive, as it leads to a wide variety of directions yet to be pursued.

References

- Allison, G. T. (1971), *Essence of decision: Explaining the Cuban missile crisis*, number 327.5 (729.1), Little, Brown and Company. [1](#)
- Arrow, K. J., Raynaud, H. et al. (1986), 'Social choice and multicriterion decision-making', *MIT Press Books* **1**. [3](#)
- Barberá, S. & Pattanaik, P. K. (1986), 'Falmagne and the rationalizability of stochastic choices in terms of random orderings', *Econometrica: Journal of the Econometric Society* pp. 707–715. [2](#), [8](#)
- Barberá, S. & Sonnenschein, H. (1978), 'Preference aggregation with randomized social orderings', *Journal of Economic Theory* **18**(2), 244–254. [3](#)
- Block, H. & Marschak, J. (1960), *Contributions to Probability and Statistics*, Stanford University Press, Stanford, CA. [2](#)
- Bossert, W. & Sprumont, Y. (2013), 'Every choice function is backwards-induction rationalizable', *Econometrica* **81**(6), 2521–2534. [3](#)
- Brualdi, R. A. & Ryser, H. J. (1991), *Combinatorial Matrix Theory*, Vol. 39, Cambridge University Press. [8](#), [13](#)
- Falmagne, J.-C. (1978), 'A representation theorem for finite random scale systems', *Journal of Mathematical Psychology* **18**(1), 52–72. [2](#), [8](#), [9](#)
- Fiorini, S. (2004), 'A short proof of a theorem of Falmagne', *Journal of Mathematical Psychology* **48**(1), 80–82. [8](#), [15](#)
- Fishburn, P. C. (1978), 'Choice probabilities and choice functions', *Journal of Mathematical Psychology* **18**(3), 205–219. [5](#)
- Ford Jr, L. R. & Fulkerson, D. R. (2015), *Flows in networks*, Princeton University Press. [2](#), [8](#), [13](#)
- Franklin, B. (1887), *The Complete Works of Benjamin Franklin: 1772-1775*, Vol. 5, GP Putnam's Sons. [1](#)
- Gelb, L. H. & Betts, R. K. (2016), *The irony of Vietnam: The system worked*, Brookings Institution Press. [1](#)
- Glass, B. (1988), *The Correspondence of Charles Darwin. Volume 2: 1837-1843*, University of Chicago Press. [1](#)
- Hammond, J. S., Keeney, R. L. & Raiffa, H. (1998), 'Even swaps: A rational method for making trade-offs', *Harvard business review* **76**, 137–150. [3](#)
- Heller, I. & Tompkins, C. (1956), 'An extension of a theorem of Dantzig's', *Linear inequalities and related systems* **38**, 247–252. [4](#), [20](#)
- Hoffman, A. J. & Kruskal, J. B. (2010), Integral boundary points of convex polyhedra, in '50 Years of Integer Programming 1958-2008', Springer, pp. 49–76. [4](#), [20](#)
- Huber, J. & Puto, C. (1983), 'Market boundaries and product choice: Illustrating attraction and substitution effects', *The Journal of Consumer Research* **10**(1), 31–44. [7](#)
- Intriligator, M. D. (1973), 'A probabilistic model of social choice', *The Review of Economic Studies* **40**(4), 553–560. [3](#)
- Intriligator, M. D. (1982), 'Probabilistic models of choice', *Mathematical Social Sciences* **2**(2), 157–166. [3](#)
- Kalai, G., Rubinstein, A. & Spiegel, R. (2002), 'Rationalizing choice functions by multiple rationales', *Econometrica* **70**(6), 2481–2488. [3](#)

- Knoblauch, V. (2016), 'Elections generate all binary relations on infinite sets', *Mathematical Social Sciences* **84**, 105–108. [6](#)
- Köksalan, M. M., Wallenius, J. & Zionts, S. (2011), *Multiple criteria decision making: from early history to the 21st century*, World Scientific. [1](#), [3](#)
- Luce, R. D. & Raiffa, H. (1957), 'Games and decisions: Introduction and critical survey (New york, 1957)', *Chs. vi and xiv* . [4](#)
- Manzini, P. & Mariotti, M. (2014), 'Stochastic choice and consideration sets', *Econometrica* **82**(3), 1153–1176. [3](#)
- Manzini, P., Mariotti, M. & Tyson, C. J. (2013), 'Two-stage threshold representations', *Theoretical Economics* **8**(3), 875–882. [5](#)
- McFadden, D. (1978), 'Modeling the choice of residential location', *Transportation Research Record* (673). [2](#)
- McGarvey, D. C. (1953), 'A theorem on the construction of voting paradoxes', *Econometrica: Journal of the Econometric Society* pp. 608–610. [2](#), [6](#)
- Pattanaik, P. K. & Peleg, B. (1986), 'Distribution of power under stochastic social choice rules', *Econometrica: Journal of the Econometric Society* pp. 909–921. [3](#)
- Payne, J. W. & Puto, C. (1982), 'Adding asymmetrically dominated alternatives: Violations of regularity and the similarity hypothesis', *Journal of Consumer Research* **9**(1), 90–98. [7](#)
- Rieskamp, J., Busemeyer, J. R. & Mellers, B. A. (2006), 'Extending the bounds of rationality: evidence and theories of preferential choice', *Journal of Economic Literature* pp. 631–661. [7](#)
- Saito, K. (2017), 'Axiomatizations of the mixed logit model', *mimeo* . [3](#), [8](#), [9](#)
- Shafir, E. (1993), 'Choosing versus rejecting: Why some options are both better and worse than others', *Memory & cognition* **21**(4), 546–556. [2](#)
- Shafir, E., Simonson, I. & Tversky, A. (1993), 'Reason-based choice', *Cognition* **49**(1), 11–36. [1](#)
- Stanley, R. P. (1997), *Enumerative Combinatorics. Vol. 1, vol. 49 of Cambridge Studies in Advanced Mathematics*, Cambridge University Press. [15](#)
- Stearns, R. (1959), 'The voting problem', *The American Mathematical Monthly* **66**(9), 761–763. [6](#)
- Telhami, S. (1990), 'Israeli foreign policy: A static strategy in a changing world', *Middle East Journal* **44**(3), 399–416. [1](#)
- Tversky, A. (1972), 'Elimination by aspects: A theory of choice.', *Psychological review* **79**(4), 281. [7](#)

5 Appendix

5.1 Proof of Theorem 3

We start by proving some lemmas that are critical for proving the theorem. First, we report the original Theorem of [Ford Jr & Fulkerson \(2015\)](#).²² Then, we prove Theorem 3, which offers a generalization of the result to any real-valued row and column vectors.

Theorem 4 (Ford Jr & Fulkerson (2015)) *Let $R = [r_1, \dots, r_m]$ and $C = [c_1, \dots, c_n]$ be positive real-valued vectors with $\sum_{i=1}^m r_i = \sum_{j=1}^n c_j$. There is an $m \times n$ matrix $A = [a_{ij}]$ such that A has row sum vector R and column sum vector C , and each entry $a_{ij} \in [0, 1]$ if and only if for each $I \subset \{1, 2, \dots, m\}$ and $J \subset \{1, 2, \dots, n\}$,*

$$|I||J| \geq \sum_{i \in I} r_i - \sum_{j \notin J} c_j. \quad (\text{FF})$$

Lemma 1 *Let $R = [r_1, \dots, r_m]$ and $C = [c_1, \dots, c_n]$ be positive real-valued vectors with $0 \leq r_i \leq 1$ and $0 \leq c_j \leq m$ such that $\sum_{i=1}^m r_i = \sum_{j=1}^n c_j$. Then there is an $m \times n$ matrix $A = [a_{ij}]$ such that A has row sum vector R and column sum vector C , and each entry $a_{ij} \in [0, 1]$.*

Proof. Given such R and C , since for each $i \in \{1, 2, \dots, m\}$, $0 \leq r_i \leq 1$, we have for each $I \subset \{1, 2, \dots, m\}$, $\sum_{i \in I} r_i \leq |I|$. Then, it directly follows that (FF) holds and conclusion follows from Theorem 4. ■

By using Lemma 1, we prove Theorem 3 that is formulated and discussed in Section 3.2.

Proof of Theorem 3 (Generalized Ford-Fulkerson Theorem). Since r_i and c_j values can be positive or negative, although the sum of the rows equals the sum of the column, their absolute values may not be the same. We analyze two cases separately, where $\sum_{i=1}^m |r_i| \geq \sum_{j=1}^n |c_j|$ and $\sum_{i=1}^m |r_i| < \sum_{j=1}^n |c_j|$. Before proceeding with these cases, first we introduce some notation and make some elementary observations.

For each real number x , let $x^+ = \max\{x, 0\}$ and $x^- = \min\{x, 0\}$. Note that for each x , $x^+ + x^- = x$. Let $R^+ = [r_1^+, \dots, r_m^+]$ and $R^- = [r_1^-, \dots, r_m^-]$. Define the n -vectors C^+ and C^- respectively. Next, let $\Sigma_{R^+} = \sum_{i=1}^m r_i^+$, $\Sigma_{R^-} = \sum_{i=1}^m r_i^-$, $\Sigma_{C^+} = \sum_{j=1}^n c_j^+$ and $\Sigma_{C^-} = \sum_{j=1}^n c_j^-$. That is, Σ_{R^+} (Σ_{R^-}) and Σ_{C^+} (Σ_{C^-}) are the sum of the positive (negative) rows in R and columns in C . Since the sum of the rows equals the sum of the columns, we have $\Sigma_{R^+} + \Sigma_{R^-} = \Sigma_{C^+} + \Sigma_{C^-}$.

For each row vector R and column vector C , suppose for each $i \in \{1, \dots, m_1\}$, $r_i \geq 0$ and for each $i \in \{m_1 + 1, \dots, m\}$, $r_i < 0$. Similarly, suppose for each $j \in \{1, \dots, n_1\}$, $c_j \geq 0$ and for each $j \in \{n_1 + 1, \dots, n\}$, $c_j < 0$. Now, let $R^1(R^2)$ be the m_1 -vector ($(m - m_1)$ -vector), consisting of the non-negative (negative) components of R . Similarly, for each column vector C , let $C^1(C^2)$ be the n_1 -vector ($(n - n_1)$ -vector), consisting of the non-negative (negative) components of C . It directly follows from the definitions that $\sum_{i=1}^{m_1} r_i = \sum_{i=1}^{m_1} r_i^+$ and $\sum_{i=m_1+1}^m r_i = \sum_{i=1}^{m_1} r_i^-$. Similarly, $\sum_{j=1}^{n_1} c_j = \sum_{j=1}^{n_1} c_j^+$ and $\sum_{j=n_1+1}^n c_j = \sum_{j=1}^{n_1} c_j^-$.

Case 1: Suppose that $\sum_{i \in I} |r_i| \geq \sum_{j \in J} |c_j|$ and let

$$\delta = \frac{\Sigma_{R^+} - \Sigma_{C^+}}{n}.$$

Note that since $\sum_{i=1}^m |r_i| \geq \sum_{j=1}^n |c_j|$, we have $\Sigma_{R^+} \geq \Sigma_{C^+}$ and $\Sigma_{R^-} \leq \Sigma_{C^-}$. Moreover, since the sum of the rows equals the sum of the columns, we have $\Sigma_{R^+} - \Sigma_{C^+} = \Sigma_{C^-} - \Sigma_{R^-}$. Therefore, by the choice of δ , we get

$$\sum_{i=1}^{m_1} r_i^+ = \sum_{j=1}^{n_1} c_j^+ + \delta \quad \text{and} \quad \sum_{i=1}^{m_1} r_i^- = \sum_{j=1}^{n_1} c_j^- - \delta. \quad (5)$$

²²This result, as stated in Lemma 4, but with integrality assumptions on R , C , and A follows from Corollary 1.4.2 in [Brualdi & Ryser \(1991\)](#). They report that [Ford Jr & Fulkerson \(2015\)](#) prove, by using network flow techniques, that the theorem remains true if the integrality assumptions are dropped, and the conclusion asserts the existence of a real nonnegative matrix.

Next, consider row-column vector pairs $(R^1, C^+ + \epsilon)$ and $(-R^2, -(C^- - \epsilon))$, where ϵ is the non-negative n -vector such that for each $j \in \{1, \dots, n\}$, $\epsilon_j = \delta$. It follows from (5) that for both pairs the sum of the rows equals the sum of the columns. Now we apply Lemma 1 to the row-column vector pairs $(R^1, C^+ + \epsilon)$ and $(-R^2, -(C^- - \epsilon))$. It directly follows that there exists a positive $m_1 \times n$ matrix A^+ and a negative $(m - m_1) \times n$ matrix A^- that satisfy (i) and (ii). We will obtain the desired matrix A by augmenting A^+ and A^- . We illustrate A^+ and A^- below.

	$(c_1^+ + \epsilon_1)$	$(c_2^+ + \epsilon_2)$	$(c_3^+ + \epsilon_3)$	\cdots	$(c_n^+ + \epsilon_n)$	
$r_1 \geq 0$	A^+					
$r_2 \geq 0$						
\vdots						
$r_{m_1} \geq 0$						
	A^-					$r_{m_1+1} < 0$
						\vdots
						$r_m < 0$
	$(c_1^- - \epsilon_1)$	$(c_2^- - \epsilon_2)$	$(c_3^- - \epsilon_3)$	\cdots	$(c_n^- - \epsilon_n)$	

Since A^+ and A^- satisfy (i) and (ii), A satisfies (i) and (ii). To see that A satisfies (iii), for each $j \in \{1, \dots, n\}$, consider $\sum_{i=1}^m |a_{ij}|$. Note that, by the construction of A^+ and A^- , for each $j \in \{1, \dots, n\}$,

$$\sum_{i=1}^m |a_{ij}| = c_j^+ + \epsilon_j + (-c_j^- + \epsilon_j) = |c_j| + 2\epsilon_j = |c_j| + 2 \frac{\Sigma_{R^+} - \Sigma_{C^+}}{n}. \quad (6)$$

Since for each $j \in \{1, \dots, n\}$, $c_j = c_j^+ + c_j^-$ such that either $c_j^+ = 0$ or $c_j^- = 0$, we get $|c_j| = c_j^+ - c_j^-$. To see that (iii) holds, observe that $\sum_{i=1}^m |r_i| - \sum_{j=1}^n |c_j| = \Sigma_{R^+} - \Sigma_{C^+} + \Sigma_{C^-} - \Sigma_{R^-}$. Since the sum of the rows equals the sum of the columns, i.e. $\Sigma_{R^+} + \Sigma_{R^-} = \Sigma_{C^+} + \Sigma_{C^-}$, we also have $\Sigma_{R^+} - \Sigma_{C^+} = \Sigma_{C^-} - \Sigma_{R^-}$. This observation, together with (6), implies that (iii) holds.

Case 2 Suppose that $\sum_{i=1}^m |r_i| < \sum_{j=1}^n |c_j|$. First, we show that there exists a non-negative m -vector ϵ such that

(E1) for each $i \in \{1, \dots, m\}$, $r_i^+ + \epsilon_i \leq 1$ and $r_i^- - \epsilon_i \geq -1$, and

(E2) $\sum_{i=1}^m r_i^+ + \epsilon_i = \sum_{j=1}^n c_j^+$ (equivalently $\sum_{i=1}^m r_i^- - \epsilon_i = \sum_{j=1}^n c_j^-$) holds.

Step 1: We show that if $\Sigma_{C^+} - \Sigma_{R^+} \leq m - \sum_{i=1}^m |r_i|$, then there exists a non-negative m -vector ϵ that satisfies (E1) and (E2). To see this, first note that $m - \sum_{i=1}^m |r_i| = \sum_{i=1}^m (1 - |r_i|)$. Next, note that, by simply rearranging the terms, we can rewrite (E2) as follows:

$$\sum_{i=1}^m \epsilon_i = \Sigma_{C^+} - \Sigma_{R^+}. \quad (7)$$

Since $\Sigma_{C^+} - \Sigma_{R^+} \leq \sum_{i=1}^m (1 - |r_i|)$, for each $i \in \{1, \dots, m\}$, we can choose an ϵ_i such that $0 \leq \epsilon_i \leq 1 - |r_i|$ and (7) holds. It directly follows that the associated ϵ vector satisfies (E1) and (E2).

Step 2: We show that since $2m \geq \sum_{i=1}^m |r_i| + \sum_{j=1}^n |c_j|$, we have $\Sigma_{C^+} - \Sigma_{R^+} \leq m - \sum_{i=1}^m |r_i|$. First, it directly follows from the definitions that

$$\sum_{i=1}^m |r_i| + \sum_{j=1}^n |c_j| = \Sigma_{R^+} - \Sigma_{R^-} + \Sigma_{C^+} - \Sigma_{C^-}.$$

Since the sum of the rows equals the sum of the columns, i.e. $\Sigma_{R^+} + \Sigma_{R^-} = \Sigma_{C^+} + \Sigma_{C^-}$, we also have $\Sigma_{R^+} - \Sigma_{C^-} = \Sigma_{C^+} - \Sigma_{R^-}$. It follows that

$$\Sigma_{C^+} - \Sigma_{R^-} \leq m.$$

Finally, if we subtract $\sum_{i=1}^m |r_i|$ from both sides of this equality, we obtain $\Sigma_{C^+} - \Sigma_{R^+} \leq m - \sum_{i=1}^m |r_i|$, as desired.

Proof. First, note that (10) can be written as follows:

$$\sum_{a \in S} |q(a, S)| + \sum_{b \notin S} |-q(b, S)| \leq 2^k. \quad (11)$$

For a set of real numbers, $\{x_1, x_2, \dots, x_n\}$, to show $\sum_{i=1}^n |x_i| \leq 2d$, it suffices to show that for each $I \subset \{1, 2, \dots, n\}$, we have $-d \leq \sum_{i \in I} x_i \leq d$. Now, as the set of real numbers, consider $\{q(a, S)\}_{a \in X}$. It follows that to show that (11) holds, it suffices to show that for each $S_1 \subset S$ and $S_2 \subset X \setminus S$,

$$-2^{k-1} \leq \sum_{a \in S_1} q(a, S) - \sum_{b \in S_2} q(b, S) \leq 2^{k-1}$$

holds. To see this, first, for each $S_1 \subset S$ and $S_2 \subset X \setminus S$, it follows from Lemma 2 that for each $a \in S_1$ and for each $b \in S_2$, we have

$$q(a, S) = \sum_{S \subset T \subset X} (-1)^{|T|-|S|} p(a, T) \quad \text{and} \quad q(b, S) = \sum_{S \subset T \subset X} (-1)^{|T|-|S|-1} p(b, T). \quad (12)$$

Note that we obtain the second equality from Lemma 2, since for each $b \notin S$, by definition of $q(b, S)$, we have $q(b, S) = q(b, S \cup \{b\})$. Next, note that for each $T \in \Omega$ with $S \subset T$, $a \in S$, and $b \notin S$, $p(a, T)$ has the opposite sign of $p(b, T)$. Now, suppose for each $b \in S_2$, we multiply $q(b, S)$ with -1 . Then, it follows from (12) that

$$\sum_{a \in S_1} q(a, S) - \sum_{b \in S_2} q(b, S) = \sum_{S \subset T \subset X} (-1)^{|T|-|S|} \sum_{a \in S_1 \cup S_2} p(a, T). \quad (13)$$

Note that, for each $T \in \Omega$ such that $S \subset T$, $\sum_{a \in S_1 \cup S_2} p(a, T) \in [0, 1]$. Therefore, the term $(-1)^{|T|-|S|} \sum_{a \in S_1 \cup S_2} p(a, T)$ adds at most 1 to the right-hand side of (13) if $|T| - |S|$ is even, and at least -1 if $|T| - |S|$ is odd. Since $|S| = n - k$, for each m with $n - k \leq m \leq n$, there are $\binom{k}{m-n+k}$ possible choice sets $T \in \Omega$ such that $S \subset T$ and $|T| = m$. Moreover, for each $i \in \{1, \dots, k\}$, there are $\binom{k}{i}$ possible choice sets T such that $S \subset T$ and $|T| = n - k + i$. Now, the right-hand side of (13) reaches its maximum (minimum) when the negative (positive) terms are 0 and the positive (negative) terms are $1(-1)$. Thus, we get

$$-\sum_{i=0}^{\lfloor \frac{k-1}{2} \rfloor} \binom{k}{2i+1} \leq \sum_{S \subset T \subset X} (-1)^{|T|-|S|} \sum_{a \in S_1 \cup S_2} p(a, T) \leq \sum_{i=0}^{\lfloor \frac{k}{2} \rfloor} \binom{k}{2i}.$$

It follows from the *binomial theorem* that both leftmost and rightmost sums are equal to 2^{k-1} . This, combined with (13), implies

$$-2^{k-1} \leq \sum_{a \in S_1} q(a, S) - \sum_{b \in S_2} q(b, S) \leq 2^{k-1}.$$

Then, as argued before, it follows that $\sum_{a \in X} |q(a, S)| \leq 2^k$. ■

Now, we are ready to complete the proof of Theorem 2. Recall that we assume $|X| = n$. For each $k \in \{1, \dots, n\}$, let $\Omega_k = \{S \in \Omega : |S| > n - k\}$. Note that $\Omega_n = \Omega$ and $\Omega_1 \subset \Omega_2 \subset \dots \subset \Omega_n$. For each pair of orderings $\succ_1, \succ_2 \in \mathcal{P}$, \succ_1 is k -**identical** to \succ_2 , denoted by $\succ_1 \sim_k \succ_2$, if the first k -ranked alternatives are the same. Note that \sim_k is an equivalence relation on \mathcal{P} . Let \mathcal{P}^k be the collection of orderings, such that each set (equivalence class) contains orderings that are k -identical to each other (\mathcal{P}^k is the quotient space induced from \sim_k). For each $k \in \{1, \dots, n\}$, let $[\succ^k]$ denote an **equivalence class** at \mathcal{P}^k , where \succ^k linearly orders a fixed set of k alternatives in X .

Note that for each $k \in \{1, \dots, n\}$, $S \in \Omega_k$ and $\succ_1, \succ_2 \in \mathcal{P}$, if $\succ_1 \sim_k \succ_2$, then since S contains more than $n - k$ alternatives, $\max(\succ_1, S) = \max(\succ_2, S)$. Therefore, for each $S \in \Omega_k$, it is sufficient to specify the weights on the equivalence classes contained in \mathcal{P}^k instead of all the weights over \mathcal{P} . Let p_k be the restriction of p to Ω_k . Similarly, if λ is a signed weight function over \mathcal{P} , then let λ^k be the restriction of λ to \mathcal{P}^k , i.e. for each $[\succ^k] \in \mathcal{P}^k$, $\lambda^k[\succ^k] = \sum_{\succ_i \in [\succ^k]} \lambda(\succ_i)$. It directly follows that λ

represents p if and only if for each $k \in \{1, \dots, n\}$, λ^k represents p_k . In what follows, we inductively show that for each $k \in \{1, \dots, n\}$, there is a signed weight function λ^k over \mathcal{P}^k that represents p_k . For $k = n$ we obtain the desired λ .

For $k = 1$, $\Omega_1 = \{X\}$ and \mathcal{P}^1 consists of n -many equivalence classes such that each class contains all the orderings that top rank the same alternative, irrespective of whether these are chosen with positive probability. That is, if $X = \{x_1, \dots, x_n\}$, then $\mathcal{P}^1 = \{[\succ^{x_1}], \dots, [\succ^{x_n}]\}$, where for each $i \in \{1, \dots, n\}$ and $\succ_i \in [\succ^{x_i}]$, $\max(X, \succ_i) = x_i$. Now, for each $x_i \in X$, define $\lambda^1([\succ^{x_i}]) = p(x_i, X)$. It directly follows that λ^1 is a signed weight function over \mathcal{P}^1 that represents p_1 .

For $k = 2$, $\Omega_2 = \{X\} \cup \{X \setminus \{x\}\}_{x \in X}$ and \mathcal{P}^2 consists of $\binom{n}{2}$ -many equivalence classes such that each class contains all the orderings that top rank the same two alternatives. Now, for each $[\succ_i^2] \in \mathcal{P}^2$ such that x_{i1} is the first-ranked alternative and x_{i2} is the second-ranked alternative, define $\lambda^2([\succ_i^2]) = p(x_{i2}, X \setminus \{x_{i1}\}) - p(x_{i2}, X)$. It directly follows that λ^2 is a signed weight function over \mathcal{P}^2 that represents p_2 . Next, by our inductive hypothesis, we assume that for each $k \in \{1, \dots, n-1\}$, there is a signed weight function λ^k over \mathcal{P}^k that represents p_k . Next, we show that we can construct λ^{k+1} over \mathcal{P}^{k+1} that represents p_{k+1} .

Note that \mathcal{P}^{k+1} is a refinement of \mathcal{P}^k , in which each equivalence class $[\succ^k] \in \mathcal{P}^k$ is divided into sub-equivalence classes $\{[\succ_1^{k+1}], \dots, [\succ_{n-k}^{k+1}]\} \subset \mathcal{P}^{k+1}$. Given λ^k , we require λ^{k+1} satisfy for each $[\succ^k] \in \mathcal{P}^k$ the following

$$\lambda^k([\succ^k]) = \sum_{j=1}^{n-k} \lambda^{k+1}([\succ_j^{k+1}]). \quad (14)$$

If λ^{k+1} satisfies (14), then since induction hypothesis implies that λ^k represents p_k , we get for each $S \in \Omega_k$ and $x \in S$, $p(x, S) = \lambda^{k+1}(\{[\succ_j] \in \mathcal{P}^{k+1} : x = \max(S, \succ_j)\})$.

Next, we show that λ^{k+1} can be constructed such that (14) holds, and for each $S \in \Omega_{k+1} \setminus \Omega_k$, λ^{k+1} represents $p_{k+1}(S)$. To see this, pick any $S \in \Omega_{k+1} \setminus \Omega_k$. It follows that $|S| = n - k$. Let $S = \{x_1, \dots, x_{n-k}\}$ and $X \setminus S = \{y_1, y_2, \dots, y_k\}$. Recall that each $[\succ^k] \in \mathcal{P}^k$ linearly orders a fixed set of k -many alternatives. Let $\{\succ^k\}$ denote the set of k alternatives ordered by \succ^k . Now, there exist $k!$ -many $[\succ^k] \in \mathcal{P}^k$ such that $\{\succ^k\} = X \setminus S$. Let $\{[\succ_1^k], \dots, [\succ_{k!}^k]\}$ be the collection of all such classes. Each ordering that belongs to one of these classes is a different ordering of the same set of k alternatives.

Now, let $I = \{1, \dots, k!\}$ and $J = \{1, \dots, n - k\}$. For each $i \in I$ and $j \in J$, suppose that \succ_{ij}^{k+1} linearly orders $X \setminus S$ as in \succ_i^k and ranks x_j in the $k+1$ th position. Consider the associated equivalence class $[\succ_{ij}^{k+1}]$. Next, we specify $\lambda^{k+1}([\succ_{ij}^{k+1}])$, the signed weight of $[\succ_{ij}^{k+1}]$, such that the resulting λ^{k+1} represents p_{k+1} . To see this, we proceed in two steps.

Step 1: First, we show that for each $S \in \Omega_{k+1} \setminus \Omega_k$, if the associated $\{\lambda_{ij}^{k+1}\}_{ij \in I \times J}$ satisfies the following two equalities for each $i \in I$ and $j \in J$,

$$\sum_{j \in J} \lambda_{ij}^{k+1} = \lambda^k([\succ_i^k]) \quad (\text{RS})$$

$$\sum_{i \in I} \lambda_{ij}^{k+1} = q(x_j, S) \quad (\text{CS})$$

then λ^{k+1} represents $p_{k+1}(S)$. For each $S \in \Omega$ and $x_j \in S$, $q(x_j, S)$ is as defined in (9) by using the given RCF p .

For each $S \in \Omega$ and $a \in S$, let $B(a, S)$ be the collection of all orderings at which a is the top-ranked alternative in S , and for each $k \in \mathbb{N}$ such that $n - k \leq |S|$, $\mathbf{B}^{k+1}(a, S)$ be the set of associated equivalence classes in \mathcal{P}^{k+1} , i.e. $B(a, S) = \{\succ \in \mathcal{P} : a = \max(S, \succ)\}$ and $\mathbf{B}^{k+1}(a, S) = \{[\succ^{k+1}] \in \mathcal{P}^{k+1} : [\succ^{k+1}] \subset B(a, S)\}$. To prove the result we have to show that for each $x_j \in S$,

$$p(x_j, S) = \sum_{\{[\succ^{k+1}] \in \mathbf{B}^{k+1}(x_j, S)\}} \lambda^{k+1}([\succ^{k+1}]). \quad (15)$$

To see this, for each $\succ \in \mathcal{P}$ and $a \in X$, let $W(\succ, a)$ denote the set of alternatives that are worse than a at \succ and a itself, i.e. $W(\succ, a) = \{x \in X : a \succ x\} \cup \{a\}$. For each $S \in \Omega$ with $a \in X$. Let $Q(a, S)$

be the collection of all orderings such that $W(\succ, a)$ is exactly $S \cup \{a\}$ and for each $k \in \mathbb{N}$ such that $n - k \leq |S|$, $\mathbf{Q}^{k+1}(a, S)$ be the set of associated equivalence classes in \mathcal{P}^{k+1} , i.e. $Q(a, S) = \{\succ \in \mathcal{P} : W(\succ, a) = S \cup \{a\}\}$ and $\mathbf{Q}^{k+1}(a, S) = \{[\succ^{k+1}] \in \mathcal{P}^{k+1} : [\succ^{k+1}] \subset Q(a, S)\}$. Note that, for each $x_j \in S$, we have $Q(x_j, S) = \bigcup_{i \in I} [\succ_{ij}^{k+1}]$. Moreover, it directly follows from the definitions of $Q(x_j, \cdot)$ and $B(x_j, \cdot)$ that

$$B(x_j, S) = \bigcup_{S \subset T} Q(x_j, T). \quad (16)$$

It follows from this observation that the right-hand side of (15) can be written as

$$\sum_{S \subset T} \sum_{\{[\succ^{k+1}] \in \mathbf{Q}^{k+1}(x_j, T)\}} \lambda^{k+1}([\succ^{k+1}]). \quad (17)$$

i. Since (CS) holds, we have

$$q(x_j, S) = \sum_{\{[\succ^{k+1}] \in \mathbf{Q}^{k+1}(x_j, S)\}} \lambda^{k+1}([\succ^{k+1}]). \quad (18)$$

ii. Next, we argue that for each $T \in \Omega$ such that $S \subsetneq T$,

$$q(x_j, T) = \sum_{\{[\succ^{k+1}] \in \mathbf{Q}^{k+1}(x_j, T)\}} \lambda^{k+1}([\succ^{k+1}]). \quad (19)$$

To see this, recall that by definition of $q(x_j, T)$ (9), we have

$$q(x_j, T) = \sum_{T \subset T'} (-1)^{|T'| - |T|} p(x_j, T'). \quad (20)$$

Since by the induction hypothesis, λ^k represents p_k , we have

$$p(x_j, T') = \sum_{\{[\succ^k] \in \mathbf{B}^k(x_j, T')\}} \lambda^k([\succ^k]). \quad (21)$$

Next, suppose that we substitute (21) into (20). Now, consider the set collection $\{B(x_j, T')\}_{\{T \subset T'\}}$. Note that if we apply the *principle of inclusion-exclusion* to this set collection, then we obtain $Q(x_j, T)$. It follows that

$$\sum_{T \subset T'} (-1)^{|T'| - |T|} \sum_{\{[\succ^k] \in \mathbf{B}^k(x_j, T')\}} \lambda^k([\succ^k]) = \sum_{\{[\succ^k] \in \mathbf{Q}^k(x_j, T)\}} \lambda^k([\succ^k]). \quad (22)$$

Since (RS) holds, we have

$$\sum_{\{[\succ^k] \in \mathbf{Q}^k(x_j, T)\}} \lambda^k([\succ^k]) = \sum_{\{[\succ^{k+1}] \in \mathbf{Q}^{k+1}(x_j, T)\}} \lambda^{k+1}([\succ^{k+1}]). \quad (23)$$

Thus, if we combine (20)-(23), then we obtain that (19) holds.

Now, (17) combined with (18) and (19) imply that the right-hand side of (15) equals to $\sum_{S \subset T} q(x_j, T)$. Finally, it follows from Lemma 2 that

$$p(x_j, S) = \sum_{S \subset T} q(x_j, T). \quad (24)$$

Thus, we obtain that (15) holds.

In what follows we show that for each $S \in \Omega_{k+1} \setminus \Omega_k$, there exists $k! \times (n - k)$ matrix $\lambda = [\lambda_{ij}^{k+1}]$ such that both (RS) and (CS) holds, and each $\lambda_{ij}^{k+1} \in [-1, 1]$. To prove this we use Theorem 3. For this, for each $i \in I$ let $r_i = \lambda^k([\succ_i^k])$ and for each $j \in J$ let $c_j = q(x_j, S)$. Then, let $R = [r_1, \dots, r_{k!}]$ and $C = [c_1, \dots, c_{n-k}]$. In Step 2, we show that the sum of C equals the sum of R . In Step 3, we show that for each $k > 1$, $2k! \geq \sum_{i=1}^{k!} |r_i| + \sum_{j=1}^{n-k} |c_j|$.

Step 2: We show that the sum of C equals the sum of R , i.e.

$$\sum_{j \in J} q(x_j, S) = \sum_{i \in I} \lambda^k([\succ_i^k]). \quad (25)$$

First, if we substitute (9) for each $q(x_j, S)$, then we get

$$\sum_{j \in J} q(x_j, S) = 1 + \sum_{j \in J} \sum_{S \subsetneq T} (-1)^{|T|-|S|} p(x_j, T). \quad (26)$$

Now, let $F(x_j)$ be the collection of orderings \succ such that there exists $T \in \Omega$ such that $S \subsetneq T$ and x_j is the \succ -top-ranked alternative in T , i.e. $F(x_j) = \{\succ \in \mathcal{P} : \max(T, \succ) = x_j \text{ for some } S \subsetneq T\}$. For each $k \in \mathbb{N}$ such that $n - k \leq |S|$, let $\mathbf{F}(x_j)$ be the set of associated equivalence classes in \mathcal{P}^k . Next, we show that for each $x_j \in S$,

$$\sum_{S \subsetneq T} (-1)^{|T|-|S|+1} p(x_j, T) = \sum_{\{[\succ^k] \in \mathbf{F}(x_j)\}} \lambda^k([\succ^k]). \quad (27)$$

To see this, first, since by the induction hypothesis, λ^k represents p_k , we can replace each $p(x_j, T)$ with $\sum_{\{[\succ^k] \in \mathbf{B}^k(x_j, T)\}} \lambda^k([\succ^k])$. Next, consider the set collection $\{B(x_j, T)\}_{\{S \subsetneq T\}}$. Since $\cup_{\{S \subsetneq T\}} B(x_j, T) = F(x_j)$, it follows from the *principle of inclusion-exclusion* that (27) holds. Next, when we substitute (27) in (26), we obtain

$$\sum_{j \in J} q(x_j, S) = 1 - \sum_{\{[\succ^k] \in \mathbf{F}(x_j)\}} \lambda^k([\succ^k]). \quad (28)$$

Then, since, by the induction hypothesis, λ^k represents p_k , we can replace 1 with $\sum_{\{[\succ^k] \in \mathcal{P}^k\}} \lambda^k([\succ^k])$. Finally, note that an equivalence class $[\succ^k] \notin \cup_{j \in J} \mathbf{F}(x_j)$ if and only if $\{\succ^k\} \cap S = \emptyset$. This means $\mathcal{P}^k \setminus \cup_{j \in J} \mathbf{F}(x_j) = \{[\succ_i^k]\}_{i \in I}$. It follows that (25) holds.

Step 3: To show that the base of induction holds, we showed that for $k = 1$ and $k = 2$, the desired signed weight functions exist. To get the desired signed weight functions for each $k + 1 > 2$, we will apply Theorem 3. To apply Theorem 3, we have to show that for each $k \geq 2$, $\sum_{i=1}^k |r_i| + \sum_{j=1}^{n-k} |c_j| \leq 2k!$. In what follows we show that this is true. That is, we show that for each $S \in \Omega_{k+1} \setminus \Omega_k$

$$\sum_{i \in I} |\lambda^k([\succ_i^k])| + \sum_{j \in J} |q(x_j, S)| \leq 2k!. \quad (29)$$

To see this, first we will bound the term $\sum_{i \in I} |\lambda^k([\succ_i^k])|$. As noted before, each $i \in I = \{1, \dots, k!\}$ corresponds to a specific linear ordering of $X \setminus S$. For each $y \notin S$, there are $k - 1!$ such different orderings that rank y at the k^{th} position. So, there are $k - 1!$ different equivalence classes in \mathcal{P}^k that rank y at the k^{th} position. Let $I(y)$ be the index set of these equivalence classes. Since $\{I(y)\}_{y \notin S}$ partitions I , we have

$$\sum_{i \in I} |\lambda^k([\succ_i^k])| = \sum_{y \notin S} \sum_{i \in I(y)} |\lambda^k([\succ_i^k])|. \quad (30)$$

Now, fix $y \notin S$ and let $T = S \cup \{y\}$. Since for each $i \in I(y)$, $[\succ_i^k] \in \mathbf{Q}^k(y, T)$ and vice versa, we have

$$\sum_{i \in I(y)} |\lambda^k([\succ_i^k])| = \sum_{[\succ_i^k] \in \mathbf{Q}^k(y, T)} |\lambda^k([\succ_i^k])|. \quad (31)$$

Recall that by the definition of $q(y, T)$, we have

$$q(y, T) = \sum_{[\succ_i^k] \in \mathbf{Q}^k(y, T)} \lambda^k([\succ_i^k]). \quad (32)$$

Next, consider the construction of the values $\{\lambda^k([\succ_i^k])\}_{i \in I(y)}$ from the previous step. For $k = 2$, as indicated in showing the base of induction, there is only one row; that is, there is a single $\{[\succ_i^k]\} =$

$\mathbf{Q}^k(y, T)$. Therefore, we directly have $|\lambda^k([\succ_i^k])| = |q(y, T)|$. For $k > 2$, we construct λ^k by applying Theorem 3. It follows from iii of Theorem 3 that

$$\sum_{[\succ_i^k] \in \mathbf{Q}^k(y, T)} |\lambda^k([\succ_i^k])| \leq |q(y, T)| + \frac{(k-1)!}{n-k+1}. \quad (33)$$

Now, if we sum (33) over $y \notin S$, we get

$$\sum_{y \notin S} \sum_{[\succ_i^k] \in \mathbf{Q}^k(y, S \cup y)} |\lambda^k([\succ_i^k])| \leq \left(\sum_{y \notin S} |q(y, S \cup y)| \right) + \frac{k!}{n-k+1}. \quad (34)$$

Recall that by definition, we have $\mathbf{Q}^k(y, S \cup y) = \mathbf{Q}^k(y, S)$ and $q(y, S \cup y) = q(y, S)$. Similarly, since each $j \in J = \{1, \dots, n\}$ denotes an alternative $x_j \in S$, we have $\sum_{x \in S} |q(x, S)| = \sum_{j \in J} |q(x_j, S)|$. Now, if we add $\sum_{j \in J} |q(x_j, S)|$ to both sides of (34), then we get

$$\sum_{i \in I} |\lambda^k([\succ_i^k])| + \sum_{j \in J} |q(x_j, S)| \leq \sum_{x \in X} |q(x, S)| + \frac{k!}{n-k+1}. \quad (35)$$

Since by Lemma 3, $\sum_{x \in X} |q(x, S)| \leq 2^k$, we get

$$\sum_{i \in I} |\lambda^k([\succ_i^k])| + \sum_{j \in S} |q(x_j, S)| \leq 2^k + \frac{k!}{n-k+1}. \quad (36)$$

Finally, note that since for each k such that $2 < k < n$ $2^k \leq \frac{(2n-2k+1)k!}{n-k+1}$ holds, we have $2^k + \frac{k!}{n-k+1} \leq 2k!$. This, together with (36), implies that (29) holds. Thus, we complete the inductive construction of the desired signed weight function.

5.3 Proof of Theorem 1

We prove this result by following the construction used to prove Theorem 2. So, we proceed by induction. Note that since C is a deterministic choice function, for each $x_i \in X$, $\lambda^1([\succ^{x_i}]) \in \{0, 1\}$. Next, by proceeding inductively, we assume that for any $k \in \{1, \dots, n-1\}$, there is a signed weight function λ^k that takes values $\{-1, 0, 1\}$ over \mathcal{P}^k and represents C_k . It remains to show that we can construct λ^{k+1} taking values $\{-1, 0, 1\}$ over \mathcal{P}^{k+1} , and that represents C_{k+1} . We know from Step 1 of the proof of Theorem 2 that to show this it is sufficient to construct λ^{k+1} such that (RS) and (CS) holds. However, this time, in addition to satisfying (RS) and (CS), we require each $\lambda_{ij}^{k+1} \in \{-1, 0, 1\}$.

First, note that equalities (RS) and (CS) can be written as a system of linear equations: $A\lambda = b$, where $A = [a_{ij}]$ is a $(k! + (n-k)) \times (n-k)k!$ matrix with entries $a_{ij} \in \{0, 1\}$, and $b = [\lambda^k([\succ_1^k]), \dots, \lambda^k([\succ_{k!}^k]), q(x_1, S), \dots, q(x_{n-k}, S)]$ is the column vector of size $k! + (n-k)$. Let Q denote the associated polyhedron, i.e. $Q = \{\lambda \in \mathbb{R}^{(n-k)k!} : A\lambda = b \text{ and } -1 \leq \lambda \leq 1\}$. A matrix is **totally unimodular** if the determinant of each square submatrix is 0, 1 or -1 . Following result directly follows from Theorem 2 of Hoffman & Kruskal (2010).

Lemma 4 (Hoffman & Kruskal (2010)) *If matrix A is totally unimodular, then the vertices of Q are integer valued.*

Heller & Tompkins (1956) provide the following sufficient condition for a matrix being totally unimodular.

Lemma 5 (Heller & Tompkins (1956)) *Let A be an $m \times n$ matrix whose rows can be partitioned into two disjoint sets R_1 and R_2 . Then, A is totally unimodular if:*

1. Each entry in A is 0, 1, or -1 ;

2. Each column of A contains at most two non-zero entries;
3. If two non-zero entries in a column of A have the same sign, then the row of one is in R_1 , and the other is in R_2 ;
4. If two non-zero entries in a column of A have opposite signs, then the rows of both are in R_1 , or both in R_2 .

Next, by using Lemma 5, we show that the matrix that is used to define (RS) and (CS) as a system of linear equations is totally unimodular. To see this, let A be the matrix defining the polyhedron Q . Since $A = [a_{ij}]$ is a matrix with entries $a_{ij} \in \{0, 1\}$, (1) and (4) are directly satisfied. To see that (2) and (3) also hold, let $R_1 = [1, \dots, k!]$ consist of the first $k!$ rows and $R_2 = [1, \dots, n - k]$ consist of the remaining $n - k$ rows of A . Note that for each $i \in R_1$, the i^{th} row A_i is such that $A_i \lambda = \lambda^k \binom{k}{i}$. That is, for each $j \in \{(i - 1)k!, \dots, ik!\}$, $a_{ij} = 1$ and the rest of A_i equals 0. For each $i \in R_2$, the i^{th} row A_i is such that $A_i \lambda = q(x_i, A)$. That is, for each $j \in \{i, i + k!, \dots, i + (n - k - 1)k!\}$, $a_{ij} = 1$ and the rest of A_i equals 0. To see that (2) and (3) hold, note that for each $i, i' \in R_1$ and $i, i' \in R_2$, the non-zero entries of A_i and $A_{i'}$ are disjoint. It follows that for each column there can be at most two rows with value 1, one in R_1 and the other in R_2 .

Finally, it follows from the construction in Step 3 of the proof of Theorem 2 that Q is nonempty, since there is λ vector with entries taking values in the $[-1, 1]$ interval. Since, as shown above, A is totally unimodular, it directly follows from Lemma 4 that the vertices of Q are integer valued. Therefore, λ^{k+1} can be constructed such that (RS) and (CS) hold, and each $\lambda_{ij}^{k+1} \in \{-1, 0, 1\}$.

Serhat Dogan
 Bilkent University
 Ankara, Turkey
 Email: dserhat@bilkent.edu.tr

Kemal Yildiz
 Bilkent University
 Ankara, Turkey
 Email: kemal.yildiz@bilkent.edu.tr