

Direct interpolation for modal μ -calculus

Bahareh Afshari^{1,2} and Graham E. Leigh²

¹ University of Amsterdam, The Netherlands
b.afshari@uva.nl

² University of Gothenburg, Sweden
graham.leigh@gu.se

The modal μ -calculus is an extension of modal logic by two quantifiers, μ and ν , that bind propositional variables. The formulæ μxA and νxA are interpreted over Kripke frames (labelled transition systems) respectively as the least and greatest fixed points of the function $x \mapsto A(x)$. Modal μ -calculus can thus be thought of as a logic that allows for restricted second-order quantification while still maintaining computationally attractive properties such as decidability of validity and the finite model property. Moreover, many program logics (LTL, PDL, CTL, etc) can be embedded into μ -calculus making it an important metatheory [3, 4].

Modal logics are known to widely enjoy interpolation (see e.g. [8, 19]) and μ -calculus does so in a very strong sense: given a formula A and a finite set of propositions and modality operators L , there exists a formula B (an interpolant) in the language L such that $A \rightarrow B$ is valid and for every formula C whose common language with A lies within L , $A \rightarrow C$ is valid if and only if $B \rightarrow C$ is valid. This property, called *uniform interpolation*, easily implies *Craig interpolation*: if $A \rightarrow C$ is valid then there is a formula B in the language common to A and C such that both $A \rightarrow B$ and $B \rightarrow C$ are valid.

Uniform interpolation for modal μ -calculus was established by D'Agostino and Hollenberg in [6]. The proof is very interesting as it involves both semantic and syntactic arguments. The authors utilise (disjunctive) modal automata from [10] to show that a form of propositional quantifier, called bisimulation quantifiers ([17, 15]), is definable in modal μ -calculus and can readily be used to define the interpolants.

Aside from the method of propositional quantification, there are a number of other ways to approach interpolation in non-classical and modal logics (see e.g. [5, 12]) among which is the syntactical approach via sequent calculus. If one has at hand a (complete) sequent calculus that admits elimination of cuts then Craig interpolation is often (but not always) provable via induction over the cut-free derivations. Indeed, there is an intimate connection between interpolation and the existence of various forms of sequent calculi, as demonstrated in [9, 13].

Following the proof-theoretic approach to interpolation, in this paper we show how to directly extract interpolants for modal μ -calculus formulæ via a sequent calculus.

The first proof system to consider is Kozen's original axiomatisation [11] which expands the standard axioms of the modal system K by regeneration and induction rules for the least (μ) and greatest (ν) fixed point quantifiers:

$$\begin{array}{ll} \mu\text{-regeneration:} & A(\mu xA(x)) \rightarrow \mu xA(x) & \mu\text{-induction:} & A(B) \rightarrow B \vdash \mu xA(x) \rightarrow B \\ \nu\text{-regeneration:} & \nu xA(x) \rightarrow A(\nu xA(x)) & \nu\text{-induction:} & B \rightarrow A(B) \vdash B \rightarrow \nu xA(x) \end{array}$$

Completeness for Kozen's system was established in [18]. The proof, imitated for the natural sequent formulation of the system, makes essential use of the cut rule and it remains a significant open problem whether Kozen's system without cut is also complete. But aside from cut, the induction rules themselves do not preserve interpolants,¹ so an alternate proof calculus is still needed.

¹For example, B is an obvious interpolant for $A(B) \rightarrow B$ but it need not be an interpolant of $\mu xA \rightarrow B$.

To pursue our goal we utilise instead the finitary and complete ‘circular’ proof systems that were introduced in [1]. One of these systems discards the induction rules in favour of generalising the regeneration inferences. The new rules have the form

$$\frac{\begin{array}{c} [\Gamma \Rightarrow \Delta, \nu x A] \\ \vdots \\ \Gamma \Rightarrow \Delta, A(\nu x A) \end{array}}{\Gamma \Rightarrow \Delta, \nu x A} \nu R \qquad \frac{\begin{array}{c} [\Gamma, \mu x A \Rightarrow \Delta] \\ \vdots \\ \Gamma, A(\mu x A) \Rightarrow \Delta \end{array}}{\Gamma, \mu x A \Rightarrow \Delta} \mu L$$

where the sequent within the brackets is understood as an assumption of the proof which is discharged. Applications of the rules are subject to the condition that there is a thread from the formula $A(\sigma x A)$ in the premise to the formula $\sigma x A$ in the discharged sequent that does not regenerate fixed point variables subsuming x . This restriction is formalised by annotating formulæ: each formula in the proof is labelled by a word from a finite set of *names* in such a way as to record the regenerations of formulæ induced by the μ and ν inferences; the condition on applications of the inference above is then represented by the local requirement that the premise and discharged assumptions of the rule have identical annotations (for details, see [1]).

Our construction of interpolants from valid implications rests on utilising circular proofs which are both cut-free and analytic. The impredicativity inherent in the sequent presentation of Kozen’s axiomatisation still remains, but instead of existing at the level of inference rules, where it naturally blocks a step-wise construction of interpolants, it manifests at the level of proofs in the choice of the assumption and discharge rules utilised. Indeed, it is precisely the two rules νR and μL that introduce quantifiers into interpolants. A similar approach using circular proofs for Gödel-Löb logic GL is [16].

The interpolants obtained in this way are structurally identical to the proof witnessing the interpolated implication and from this observation one can immediately infer results on the logical form of the interpolant. For example, given a valid formula $A \rightarrow C$ there exists a conjunction-free interpolant if there is a proof of $A \rightarrow C$ avoiding the rules of conjunction introduction on the right or disjunction introduction on the left. As it is the quantifiers that are the source of expressibility and complexity in the μ -calculus, it is of interest to examine the quantifier structure of interpolants and their dependence on the structure of proofs. We classify sufficient conditions for ensuring interpolants from complexity classes Π_1 , Σ_1 and the alternation-free fragment of μ -calculus.

This work can be placed within a broader programme of adapting and applying proof-theoretic methods to coalgebraic logics in general. Modal μ -calculus is a special case of the coalgebraic fixed point logic which is known to have uniform interpolation [7, 14]. Potential avenues of future research include applying the above methods and systems to other modal and temporal logics with fixed points and a proof-theoretic treatment of uniform interpolation for μ -calculus, such as is carried out for modal logic in [2]. The former direction depends on sound and complete cut-free sequent calculi for different modal logics, a question which warrants further study independently.

References

- [1] Bahareh Afshari and Graham E. Leigh. Cut-free completeness for modal mu-calculus. In *Proceeding of Thirty-Second Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, Lecture Notes in Computer Science. Springer, 2017.
- [2] Marta Bílková. Uniform interpolation and propositional quantifiers in modal logics. *Studia Logica*, 85(1):1–31, 2007.

- [3] Julian Bradfield and Colin Stirling. Modal μ -calculi. In Patrick Blackburn, Johan van Benthem, and Frank Wolter, editors, *Handbook of Modal Logic*, pages 721–756. Elsevier, 2007.
- [4] Julian Bradfield and Igor Walukiewicz. The μ -calculus and model-checking. In Helmut Veith Edmund Clarke, Thomas Henzinger, editor, *Handbook of Model Checking*. Springer-Verlag, 2018.
- [5] Giovanna D’Agostino. Interpolation in non-classical logics. *Synthese*, 164(3):421–435, 2008.
- [6] Giovanna D’Agostino and Marco Hollenberg. Logical questions concerning the μ -calculus. *The Journal of Symbolic Logic*, 65(1):310–332, 2000.
- [7] Sebastian Enqvist, Fatemeh Seifan, and Yde Venema. Completeness for μ -calculi: A coalgebraic approach. *Ann. Pure Appl. Logic*, 170(5):578–641, 2019.
- [8] Dov M. Gabbay and Larisa Maksimova. *Interpolation and Definability. Modal and Intuitionistic Logic*. Oxford University Press, 2005.
- [9] Rosalie Iemhoff. Uniform interpolation and sequent calculi in modal logic. *Arch. Math. Log.*, 58(1-2):155–181, 2019.
- [10] David Janin and Igor Walukiewicz. *Automata for the modal μ -calculus and related results*, pages 552–562. Springer Berlin Heidelberg, Berlin, Heidelberg, 1995.
- [11] Dexter Kozen. Results on the propositional μ -calculus. *Theoretical Computer Science*, 27:333–354, 1983.
- [12] Marcus Kracht. Modal consequence relations. In Patrick Blackburn, Johan van Benthem, and Frank Wolter, editors, *Handbook of Modal Logic*, volume 3, pages 491–545. Elsevier, 2007.
- [13] Roman Kuznets. *Proving Craig and Lyndon Interpolation Using Labelled Sequent Calculi*, pages 320–335. Springer International Publishing, Cham, 2016.
- [14] Johannes Marti, Fatemeh Seifan, and Yde Venema. Uniform interpolation for coalgebraic fixpoint logic. In Lawrence S. Moss and Pawel Sobocinski, editors, *6th Conference on Algebra and Coalgebra in Computer Science, (CALCO)*, volume 35 of *LIPICs*, pages 238–252. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2015.
- [15] Andrew M. Pitts. On an interpretation of second order quantification in first order intuitionistic propositional logic. *The Journal of Symbolic Logic*, 57(1):33–52, 1992.
- [16] Danyar S. Shamkanov. Circular proofs for the Gödel-Löb provability logic. *Mathematical Notes*, 96(3):575–585, 2014.
- [17] Albert Visser. Bisimulations, model descriptions and propositional quantifiers. Logic Group Preprint Series No. 161, Utrecht, 1996.
- [18] Igor Walukiewicz. Completeness of Kozen’s axiomatisation of the propositional μ -calculus. *Information and Computation*, 157:142–182, 2000.
- [19] Michael Zakharyashev, Frank Wolter, and Alexander V. Chagrov. *Advanced Modal Logic*, pages 83–266. Springer Netherlands, 2001.